









## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> C12N 15/12, C07K 15/00 C12Q 1/68, G01N 35/567, 33/574	<b>A2</b>	<b>(11) International Publication Number:</b> WO 92/14821 <b>(43) International Publication Date:</b> 3 September 1992 (03.09.92)		
<b>(21) International Application Number:</b> PCT/US92/01375 <b>(22) International Filing Date:</b> 21 February 1992 (21.02.92) <b>(30) Priority data:</b> 660,412 22 February 1991 (22.02.91) US <b>(71) Applicant:</b> MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02139 (US). <b>(72) Inventors:</b> McKAY, Ronald, D., G. ; 3 Gorham Avenue, Brookline, MA 02146 (US). LENDAHL, Urban ; Fär-gargardstorget 70, S-116 43 Stockholm (SE).	<b>(74) Agents:</b> GRANAHAH, Patricia et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US). <b>(81) Designated States:</b> AT (European patent), BE (European patent), CA, CH (European patent), CS, DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent). <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>			
<b>(54) Title:</b> NESTIN EXPRESSION AS AN INDICATOR OF NEUROEPITHELIAL TUMORS  <table border="0" style="width: 100%;"><tr><td style="width: 50%; vertical-align: top;"></td><td style="width: 50%; vertical-align: top;"></td></tr></table> <p>A gene encoding a protein, nestin, whose expression distinguishes neural multipotential stem cells and brain tumor cells from the more differentiated neural cell types (e.g., neuronal, glial and muscle cells). Methods of detecting the expression of nestin as a means of diagnosing and treating brain tumours are also described.</p>				
				

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

-1-

NESTIN EXPRESSION AS AN INDICATOR OF  
NEUROEPITHELIAL TUMORS

Description

Background

05       Brain tumors are a leading cause of cancer deaths in people under the age of 35. An important concept in our understanding of brain tumors is that the neoplastic cells may arise from undifferentiated neuroectodermal cells (Rubinstein, L.J. J. Neurosurg 62: 795-805 (1985)).

10       At the gastrula stage of central nervous system (CNS) development, neuroectoderm, forming the neural plate is derived from ectoderm which has been induced by underlying mesoderm. (Spemann, H. Yale University Press 1938)). The neural plate then folds, in the process  
15       called neurulation, to form the neural tube. The cells of the neural tube are the precursors for the major differentiated cell types in the central nervous system: astrocytes, oligodendrocytes, and the various types of neuron (Sauer, F.C. J. Comp. Neurol. 62: 377-405  
20       (1935)). The cells of the neural tube may also be the precursors to neoplastic cells which form brain tumors.

      Recent success in "tagging" individual CNS precursors and their daughter cells, either through infection by a marker-bearing retrovirus or by injection of  
25       stable, specific dyes and enzymes, has shown that neurons and non-neuronal cells (i.e., glia cells) are often derived from a common precursor. In retina a common precursor for neurons and the Mueller glial cell exists

-2-

- very late in development (Holt, C.E., T.W. Bertsch et al., Neuron 1: 15-26 (1988); Turner, D.L. and C.L. Cepko, Nature 238: 131-136 (1987); Wetts, R. and S.E. Fraser, Science 239: 1142-1145 (1988); a common precursor for different neuronal and glial cell types has also been demonstrated in cortex and tectum (Luskin, M.B., A.L. Pearlman et al., Neuron 1: 635-647 (1988); Price, J. and L. Thurlow, Development 104: 473-482 (1988); Walsh, C. and C. Cepko, Science 241: 1342-1345 (1988)).
- 10 Tissue culture systems have provided additional information about the differentiation potential of CNS stem cells. Single-cell microculture of embryonic day 14 rat forebrain has produced clones containing both neurons and glia (Temple, S. Nature 340: 471-473 (1989)).
- 15 Analysis of a later precursor in the glial differentiation pathway has established specific factors influencing differentiation to oligodendrocytes or type II astrocytes (Raff, M. Science 243: 1450-1455 1989)). Cell lines established by immortalization of CNS stem cells
- 20 also reflect features of the stem cell and can differentiate along both the neuronal and glial pathways (Cepko, C.L. Ann. Rev. Neurosci. 12: 47-65 (1989); Frederiksen, K. et al., supra (1988)).
- Antibodies have also proven useful for analyzing stem cells. For example, antibodies A2B5 (Raff, M.C. supra 1989) D1.1 (Levine, J.M., L. Beasley et al., J. Neurosci 4: 820-831 (1984), Rat 401 (Hockfield, S. and R. McKay, J. Neurosci 5: 3310-3328 (1985), and antisera against the intermediate filament vimentin (Bignami, A.,
- 25 T. Raju et al Dev. Biol 91: 286-295 (1982); Federoff, S.
- 30

- 3 -

Molecular Bases of Neural Development (1985); Tapscott, S.J., G.S. Bennett et al., Dev. Biol. 86: 40-54 (1981) bind to antigens enriched in the proliferative zone of the neural tube. However, A2B5 and D1.1 recognize glycolipid epitopes and are less useful for analysis of gene expression. Vimentin, which appears transiently in brain development, is an ambiguous marker largely because of its promiscuous expression in many cultured cells as well as in a variety of developing and differentiated tissues (Traub, N. Intermediate filaments: a review. Berlin, SpringerVerlag (1985)).

The correlation between Rat 401 and proliferating cells in the developing nervous system has been examined in detail. Rat 401 was found to recognize a transient population of embryonic columnar epithelial cells and radial glial cells in many regions of the CNS (Hockfield, S. and McKay J. Neurosci. 5: 3310-3328 (1985)). The number of Rat 401 positive cells, their proliferative rate, and the developmental kinetics of Rat 401 expression relative to neuronal differentiation revealed that the immediate precursors to neurons are also Rat 401 positive (Frederiksen and McKay, J. J. Neurosci 8: 1144-1151 (1988)). Neither neurons nor glia in the adult brain express the epitope recognized by the Rat 401 antibody.

Although "tagging", tissue culture systems and antibodies have proven useful in analyzing neural precursor cells, there is still an incomplete understanding of the lineage of tumor development. For example, medulloblastoma, a common brain tumor of children, is associated predominantly with the cerebellum and brain

-4-

stem. Medulloblastoma tumors contain cell types with differentiated characteristics of neurons, glia and muscle (Rubinstein, supra 1985; Coakham H.B., et al., J. Clin. Pathol. 38: 165-173 (1985); Velasco, M.E. et al., 05 Surg. Neurol. 23: 177-182 (1985); Hayashi, K. et al., Acta Pathologica Japonica 37: 85-96 (1987); Cras, P. et al., Acta. Neuropathol 75: 377-384 (1988)).

Because of the presence of multiple differentiated cell types found in these tumors, Bailey and Cushing 10 (Bailey, P. and H. Cushing, Arch. Neurol. Psychiatry 14: 192-224 (1925) proposed that rather than being multiple coincident tumors, they were derived from a multi-potential stem cell, which they called a "medulloblast". According to this model, a medulloblastoma tumor would 15 result from uncontrolled proliferation and differentiation of the medulloblast. However, the hypothetical medulloblast has not been identified.

Due to the incomplete understanding of brain tumor development, there is currently no adequate assay for 20 detecting brain tumors at an early stage in their development. In addition, there is no specific noninvasive method of treating brain tumors. Methods of detecting and treating brain tumors would be very useful.

#### Summary of the Invention

25 The present invention relates to a gene encoding a protein whose expression distinguishes neural multi-potential stem cells and brain tumor cells from the more differentiated neural cell types (e.g. neuronal, glial and muscle cells) of the mammalian brain.

-5-

The present invention relates to the nestin gene, particularly a nestin gene of mammalian origin, and the encoded nestin protein. As described herein, two nestin genes of mammalian origin, the rat nestin gene and the human nestin gene, have been isolated and sequenced. The coding sequence of the rat nestin gene transcript is 5415 bp, which corresponds to an expected molecular weight of approximately 200 kD for the nestin protein. The relative molecular weight of the nestin protein as determined by SDS polyacrylamide gels is approximately 240 kD. The predicted amino acid sequence of the rat nestin gene product shows that nestin defines a distinct sixth class of intermediate filament protein.

The present invention also relates to the human nestin gene and the encoded human nestin protein. The coding sequence of the human nestin gene transcript is 4854 bp. The predicted amino acid sequence is 1618 amino acids in length, and shows significant sequence homology with the rat nestin sequence.

The present invention further relates to methods of detecting the expression of nestin as a means of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an adult individual. In one embodiment, DNA present in a sample from the brain of an adult individual is hybridized to a DNA probe which is complementary to all or a portion of the nestin gene. As used herein, the term the nestin gene includes the human nestin gene whose sequence is represented herein, the rat nestin gene whose sequence is also represented herein and equivalent genes from other

-6-

species, such as those which are substantially homologous to the human or the rat gene sequence. Detection of hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor. In another embodiment, cerebral spinal fluid or a serum sample from the brain of an adult individual can be stained with anti-nestin antibodies. Detection of stained cells is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor.

#### Brief Description of the Drawings

Figure 1 is the nucleotide and deduced amino acid sequences of the nestin transcript derived from the central nervous system of rat embryos. The three intron positions (912, 1038 and 1111) are indicated. The canonical poly A addition signal (5924-5929) is underlined.

Figure 2 is the nucleotide sequence of human nestin.

Figure 3 is the deduced amino acid sequence of human nestin.

#### Detailed Description of the Invention

The present invention relates to a gene of mammalian origin encoding a protein whose expression distinguishes neural multipotential stem cells and brain tumor cells from the more differentiated neural cell types (e.g., neuronal, glial and muscle cells of the adult brain).

The present invention more particularly relates to the nestin gene and the encoded proteins and their use in



-7-

diagnosing tumors of the brain, such as medulloblastomas, glioblastomas and oligodendroglioma. As described herein, the human nestin gene and the rat nestin gene have been isolated and sequenced. In addition, expression of the nestin gene and its structural features are described. The intracellular distribution of the nestin gene product, both in an embryonic rat brain cell line (ST15A) and human brain tissue, including tumor cells, using immunocytochemical and immunohistochemical techniques is also described. Finally, nestin expression in the developing human central nervous system and brain tumor samples is also described herein.

#### CLONING AND CHARACTERIZATION OF THE RAT AND HUMAN NESTIN GENE

##### 15 The Rat Nestin Gene

A cDNA library in the expression vector lambda gt11 was constructed from poly(A)<sup>+</sup> RNA from CNS of embryonic day 15 (E15) rat embryos as described in detail in Example 1. The library was screened with the monoclonal antibody Rat 401 after induction of protein expression and immunopositive plaques were identified. The cDNA insert from the most immunopositive clone, gt11.401.19, was used as probe to isolate additional clones, gt10.401:16, gt10.401:18 and gt10.401:9, in a lambda gt10 library from rat E15 CNS constructed in parallel with the gt11 library. A number of hybridizing clones were identified and sequenced and together represent 5333 base pair (bp) of transcribed sequence, including 404 bp of 3'

- 8 -

untranslated sequence and a consensus polyA addition site.

05 The remaining portion of the nestin gene was obtained from genomic sequences. Genomic Southern blots, even when hybridized at low stringency, revealed a pattern consistent with a single rat gene. A genomic rat library in the vector Charon 4A was screened with the nestin cDNA clones. Hybridizing clones were organized by restriction mapping and the region upstream of the cDNA  
10 clone gtl0.401:9 was sequenced.

An open reading frame continues for an additional 600 bp from the 5' end of the most upstream cDNA clone. This region from the genomic clone Cha.401:14 was subdivided into three different fragments with the restriction enzyme BamH1. These three fragments all hybridized  
15 to an mRNA of the correct size, 6.2 kb, when used as probes on Northern blots with rat E15 CNS RNA.

The exact location of the 5' end was determined by a combination of S1 nuclease mapping and primer extension.  
20 A probe (bp -121 to +194 relative to +1 - the subsequently defined transcription start) was generated overlapping the end of the genomic open reading frame by polymerase chain reaction between two specific primers. This PCR product protected a 194 bp fragment in S1  
25 nuclease analysis. Primer extension from the downstream primer used to generate the PCR fragment also produced a 194 bp fragment. The fact that these two experiments gave the same fragment suggests that the transcription start site has been defined.

30 Figure 1 shows the nucleotide sequence of the rat nestin gene. The total length of the nestin gene transcript is 5945 bp, excluding the poly A tail.

- 9 -

There are two potential initiation codons at positions +127 and +160. Assuming the first methionine is the initiation codon, the coding sequence is 5415 bp. This corresponds to an expected molecular weight of approximately 200 kD; the relative molecular weight as determined by migration on SDS polyacrylamide gels is approximately 240 kD (Hockfield and McKay, supra (1985)). Sequence comparison of cDNA and genomic clones reveals three introns at positions 912, 1038, and 1111 in the transcribed region.

To confirm that the cloned gene indeed represents the gene encoding the Rat401 antigen, a synthetic peptide from the C-terminal 20 amino acids was made. The extreme C-terminus was chosen because it is not included in the clone recognized by the original monoclonal antibody, and therefore represents a distinct epitope. A polyclonal antiserum against the peptide was raised in rabbits and identifies a band in Western blots and a pattern in immunostaining experiments which are indistinguishable from that recognized by the monoclonal antibody Rat 401.

#### Structural Features of the Rat Nestin Gene

The sequence of the entire transcribed region of the rat nestin gene was compared to the Genbank and EMBL databases. The only significant similarities found are to the five classes of intermediate filaments: acidic keratins (class I); basic keratins (class II); desmin, glial fibrillary acidic protein, peripherin, and vimentin (class III); the neurofilament triplet (class IV); and nuclear lamins (class V) (Steinert and Roop, 1988)).

-10-

The similarity between the nestin gene and the genes of the other five classes of intermediate filaments ranges from 16 to 29 % at the amino acid level in a 307 amino acid long region starting close to the N-terminus of the nestin gene, corresponding to the conserved alpha-helical rod or "core" domain of the intermediate filaments. This region of the predicted nestin amino acid sequence also contains a repeated hydrophobic heptad motif characteristic of intermediate filaments. The degree of amino acid similarity is comparable to that found between different classes of intermediate filament, both in degree and in location, i.e., conserved heptad-containing alpha-helical stretches within the core domain are separated by less well conserved, non-heptad spacers (Steinert, P.M. and D.R. Roop, Ann. Rev. Biochem. 57: 593-625 (1988)). The locations of the nestin gene's three introns are not conserved with respect to those of other classes of intermediate filaments.

In the regions outside the conserved rod domain, no strong similarities to other characterized genes were found. The N-terminal domain is only 11 amino acid residues long, shorter than in other intermediate filaments. The large C-terminal domain loosely resembles those of neurofilaments in that it is highly charged, bears glutamate-rich regions, and features a repeat, in this case the 11 amino acid motif S/P-L-E-E/K-E-X-Q-E-S/L-L-R (underlined residues are strongly conserved). There are approximately 35 of these repeats in the region between amino acids 512 and 1050.

The deduced amino acid sequence of nestin suggests that it is a member of the intermediate filament protein

-11-

family. However, its degree of sequence homology to other intermediate filaments in the core domain, its dissimilarity in the head and tail domains, and its different splicing pattern suggest that it defines a new class of intermediate filament protein.

#### The Human Nestin Gene

As described in detail in Example 5, the human nestin gene was isolated using low stringency DNA hybridization of a human genomic bacteriophage lambda library with a rat nestin probe.

Figure 2 shows the nucleotide sequence of the human nestin gene. The total length of the human nestin gene sequence is 4854 bp. Figure 3 shows the deduced amino acid sequence encoded by the human nestin gene which is 1618 amino acid residues in length. The alignment of the deduced amino acid sequences of human nestin with the deduced amino acid sequence of rat nestin showed greater than 75% sequence similarity between the two sequences, with greater than 60% sequence identity.

#### 20 DEVELOPMENTAL EXPRESSION OF THE RAT AND HUMAN NESTIN GENE AND ENCODED PRODUCT

##### Developmental Expression of the Rat Nestin Gene

During development, CNS stem cells differentiate into neurons and glia on a stereotyped schedule; different brain regions become post-mitotic at different times. The nestin protein was originally identified by the Rat 401 antibody, which transiently stains radial

-12-

glial cells of the neural tube in cross sections of rat embryos (Fredericksen, K. and R. McKay, J. Neurosci 5: 3310-3328 (1988); Hockfield and McKay, *supra* (1985)).

05 The distribution of staining in the neural tube at different developmental stages, combined with quantitative double-label experiments using tritiated thymidine autoradiography and FACS cell cycle analysis, showed that expression of the Rat 401 antigen was localized to proliferating CNS stem cells during embryogenesis  
10 (Frederiksen and McKay, *supra* (1988)) and U.S. Patent Applications Serial Nos. 180,548; 201,762 and 603,803 herein incorporated by reference. These experiments have been extended by analyzing the developmental expression of the nestin gene.

15 Northern blots of RNA from E15 rat CNS or from the Rat 401 positive cell line ST15A were probed with nestin cDNA inserts, a single 6.2 kb mRNA species was identified. To investigate the temporal profile of nestin RNA expression in the developing CNS, representatives of  
20 early (cerebrum) and late (cerebellum) developing regions of the brain were compared.

In cerebrum, nestin expression was found to decline from a maximum on day E16, such that only extremely low levels of RNA could be detected at day P9 and P12 and  
25 none at all in the adult cerebral cortex, even after long exposures. The loss of nestin mRNA precisely parallels the decline of Rat 401 positive stem cells in cerebrum during neurogenesis (Frederiksen and McKay, *supra* (1988)). The postnatal cerebellum expresses the nestin  
30 gene with a maximum on day P5; no expression can be

-13-

detected after day P9 . Again, this pattern closely matches the numbers of Rat 401 positive cells in developing cerebellum.

05       - Nestin is not detected by Northern blot analysis in  
adult tissue. The loss of nestin expression coincides  
with terminal differentiation of these early multi-  
potential cells. There are two important implications of  
these results. First, nestin appears to be a general  
marker for the CNS stem cell from the newly closed E11  
10       neural tube through the postnatally developing  
cerebellum. Second, nestin expression is a consequence  
of transcriptional regulation of the gene.

          The only expression of nestin observed outside  
developing CNS is traces of nestin message from day P4  
15       developing skeletal muscle. This finding is consistent  
with previous observations from cross sections of rat  
embryos, where Rat 401 staining in the developing somites  
was observed (Hockfield and McKay, supra (1985)).  
Developing nervous tissue and developing skeletal muscle  
20       may share unexpected kinship with regard to their  
potential fates.

#### Intracellular Distribution of the Nestin Gene Product in the ST15A Cell Line

25       The associations of nestin with different components  
of the cytoskeleton were analyzed by a series of double-  
label immunocytochemistry experiments. These experiments  
were carried out in the immortalized CNS precursor cell  
line ST15A which expresses, in addition to components of  
microtubules and microfilaments, nestin and the inter-

-14-

mediate filament vimentin. ST15A produces a nestin mRNA indistinguishable by Northern blot analysis from that found in E15 CNS.

05 In the first set of double-label experiments the intracellular distribution of the nestin gene product, detected by the monoclonal Rat.401 antibody and an appropriate second antibody, was compared to the distribution of actin, tubulin and vimentin. The latter two proteins were detected by polyclonal antibodies with  
10 matching second antibodies while the actin pattern in microfilaments was visualized by rhodamine-conjugated phalloidin.

The intracellular distributions of nestin, microtubules (tubulin), and microfilaments (actin) are quite  
15 different. Microtubules are more evenly distributed in the cell with no particular aggregation around the nucleus, whereas nestin is found in a fiber-like pattern, with a pronounced perinuclear accumulation. We also observe distinct differences when microfilaments and  
20 nestin are stained in the same cell: microfilaments form straight and parallel fibers, in contrast to the gently curved fibers of nestin.

Differences in nestin and vimentin staining in the same ST15A cells are much less clear. Vimentin staining  
25 is stronger than nestin in the cell periphery, and there are regions where intensity varies between the two proteins, but common features are also evident. The overall organization of the two networks is quite similar, with filament arrays radiating out from a  
30 perinuclear center. Moreover, there are regions in cells where it seems likely that nestin and vimentin colocalize.



-15-

In another set of experiments, microfilaments and microtubules were disrupted by cytochalasin B and colchicine, respectively, and the effects on nestin distribution analyzed. When ST15A cells were treated with 10 ug/ml cytochalasin B for one hour the microfilaments became highly disorganized, while the overall morphology of the nestin network was preserved. Microtubules also remained intact after the cytochalasin B treatment. Colchicine treatment (25 ug/ml) for 24 hours caused a collapse of the microtubules. The nestin network could still be identified, although the perinuclear distribution became much more prominent. A similar pattern was observed for vimentin as has been previously reported (Hynes, R.O. and A.T.Destree, Cell 13:151-163 (1978): Monteiro, M.J. and D.W. Cleveland, J. Cell Biol. 108: 579-593 (1989). Microfilaments were essentially unaffected by the colchicine treatment. The inferred relationship of nestin to intermediate filaments based on its deduced amino acid sequence is thus supported by the intracellular distribution of the nestin gene product under various conditions.

ST15A Cells have Features Expected of a Neural Precursor to Brain Tumors

"The prevailing name 'medulloblastoma' given by Bailey and Cushing (1925), is unfortunate because there is no embryonal cell that has been identified as a medulloblast." This sentence begins the discussion of medulloblastoma in a prominent text on neural tumors (Russell, D. and L.J.Rubinstein, Pathology of Tumors of

-16-

the Nervous System, Baltimore, Williams and Wilkins (1989)). The following presents evidence that a cell line derived from rat cerebellum, ST15A, has features expected of a medulloblast.

05 ST15A was derived from postnatal day 4 rat cerebellum. In the rat, as in other mammals, much of cerebellar development occurs after birth, so that there are a large number of dividing precursor cells present at the time from which this line was derived. Thus, ST15A is  
10 derived from the appropriate tissue and stage of development. It has previously been shown that ST15A cells can differentiate along neuronal and glial pathways. The following experiments, which are described in detail in Example 3, show that ST15A cells can also differentiate  
15 into muscle. This result suggests that a single neuroectodermal cell can give rise to the different cell types found in brain tumors.

The cerebellar cell line ST15A was one of several cell lines obtained by infecting primary rat cerebellar  
20 cells with a recombinant retrovirus carrying SV40 T-antigen. A temperature sensitive allele of T-antigen was used to establish these cell lines. At the permissive temperature (i.e., when the T-antigen protein is active) the cells proliferate and express the intermediate  
25 filament protein nestin.

At the non-permissive temperature the T-antigen protein is rapidly degraded and the cells differentiate into either neurons or glial cells depending on the conditions. The differentiation into neurons and glial  
30 cells is best accomplished by growing ST15A cells in

-17-

co-culture with primary cerebellar cells. Partial differentiation has also been observed when ST15A cells are cultured alone; serum-free medium promotes neuronal differentiation and medium containing fetal calf serum promotes glial differentiation.

ST15A cells also can differentiate into muscle cells. This fate was first noted when spontaneously contracting cells were observed in long term cultures of ST15A at 39°C, the non-permissive temperature for T-antigen. Horse serum is often used to support the differentiation of primary myoblasts. When ST15A cells are grown in horse serum at 39°C, they reproducibly differentiate into multi-nucleate cells which express muscle specific proteins, regenerative action potentials and spontaneously contract.

ST15A cells grown for two days at 33°C in 10% fetal calf serum (FCS) in DMEM grew in a disorganized manner; in contrast ST15A cells grown at 33°C in 10% horse serum at 39°C became aligned. The monoclonal antibody Rat 401 recognizes the 220 kD intermediate filament protein nestin. ST15A cells express nestin at both 33°C and 39°C. A second monoclonal antibody which recognizes a skeletal and cardiac muscle isotype of troponin T was reacted with the cells (Lin, J. Biol. Chem. 263: 7309 (1988)). Immunofluorescence shows that troponin T expression is induced when ST15A cells are cultured at 39°C in horse serum.

Immunoblots of proteins extracted from ST15A cells verify that anti-troponin T antibody recognizes a strong band of 38 Kd, the appropriate molecular weight, in the

-18-

cells grown at 39°C in horse serum; at 33°C no band is present. The troponin band in ST15A co-migrates with troponin extracted from rat postnatal day 4 muscle. The double band seen in both muscle cells and differentiated  
05 ST15A cells may represent the adult and embryonic isoforms of troponin T which differ by 3,000 daltons (Lin, J. Biol. Chem. 263: 7309 (1988)). These two isoforms are derived from the same primary transcript by differential splicing. Lanes carrying the same amount of  
10 protein were also probed for nestin with the monoclonal antibody Rat 401. Nestin is present at both temperatures but at elevated levels at the higher temperature. The lower bands are proteolytic fragments of nestin which are often present in protein extracts even in the presence of  
15 protease inhibitors. Nestin is also present in postnatal rat muscle.

A family of DNA binding factors has been shown to play a critical role in the differentiation of muscle. These proteins, myoD1, myogenin, myf5 and herculin/MRF4,  
20 contain DNA binding motifs of the helix-loop-helix type in a region of homology to c-myc (Wright, W.E., et al., Cell. 56: 607 (1989); Miner, J.H. and B.Wold, Proc. Nat. Acad. Sci. 87: 1089-1093 (1990); Davis, R.L. et al., Cell. 51: 987-1000 (1987); Braun, T et al., EMBO J 8:  
25 701-709 (1989); Rhodes S.J. and S.F.Konieczny, genes and dev. 3: 2050-2061 (1989). The expression of these factors at the mRNA level in ST15A cells as they differentiate into muscle was examined.

Northern blot analysis of total RNA from differentiating ST15A cells probed with the indicated sequences  
30 was performed. Each probe hybridized only to the bands

-19-

of the expected mobilities. MyoD1 is expressed in passaging ST15A cells at the permissive temperature for the T antigen and continues to be expressed when the cells are shifted to 39°C. Myogenin is not expressed  
05 under normal growth conditions but is expressed by 2 days after confluent cells are shifted to 39°C in horse serum (HS); if confluent cells are placed in HS and held at 33°C, the induction of myogenin expression is dramatically inhibited.

10 Herculin/MRF4 was found in adult skeletal muscle but was not detected in ST15A. Myf5 gave a very weak signal in ST15A cells compared to the myoblast cell line C2C12. Nestin mRNA accumulates with time at 39°C, as does the protein.  $\beta$ -actin serves as a control for the amount of  
15 RNA at 33°C and in the short time points at 39°C; the decrease in  $\beta$ -actin signal at 3 and 7 days at 39°C may be due to the changing composition of the cytoskeleton accompanying differentiation, as levels of myoD1 mRNA appear constant throughout the experiment. These results  
20 indicate the ST15A cells express myogenic transcription factors and that the induction of myogenin correlates with the differentiation into the muscle fate.

As expected of muscle cells, ST15A cells become multinucleate as they differentiate. The most striking  
25 similarity to muscle, which initiated this series of experiments, was that ST15A cells grown to confluence in FCS at 33°C and cultured for up to 15 days at 39° in 10% horse serum can contract spontaneously. The initial action potentials were generated by a series of hyper-  
30 polarizing pulses. The final hyper-polarizing pulse was followed by a train of spontaneous action potentials. Spontaneous action potentials were seen after 12 days at 39°C in 10% horse serum.

-20-

Coincident with the appearance of contracting cells, the resting membrane potential falls and it becomes easier to stimulate trains of action potentials with hyperpolarizing current pulses. ST15A cells also  
05 twitched in response to stimuli which elicited action potentials. These results show that ST15A cells differentiate into muscle, expressing myogenic transcription factors, muscle-specific cytoskeletal components and electrical excitability.

10 An interesting question raised by the properties of ST15A cells is whether primary cerebellar cells can differentiate into the muscle fate. Dissociated cells from postnatal day 5 rat cerebellum were cultured for 6 days under conditions which promote muscle different-  
15 iation of ST15A cells (39°C in the presence of 10% horse serum) and stained with monoclonal antibody against troponin T. These culture conditions promote extensive differentiation of neurons and glia. Troponin T positive  
20 muscle cells were also seen, but these cells were very rare, occurring less than one per hundred thousand cells.

#### Developmental expression of Human Nestin Gene

To characterize human nestin and determine the timing of its induction in the normal human developing  
25 CNS, studies were carried out, to examine this class VI IF protein in the normal human developing central nervous system (CNS), human brain tumor derived cell lines, and tissue samples of human CNS tumors.

Human nestin exhibited biochemical and immuno-  
30 chemical properties similar to those of rat nestin.

-21-

Further, in the human, nestin was detected immuno-histochemically in several different types of immature human CNS cells, i.e. germinal matrix cells, neuro-epithelial cells lining the central canal, radial glia and endothelial cells. Nestin appeared in these cells at the earliest gestational age (i.e., 6 weeks) examined here and then it declined in all but the endothelial cells at later embryonic stages. Nestin also was detected by immunocytochemistry in 6 of 7 primitive neuroectodermal tumor cell lines and in both of the malignant glioma cell lines examined. In these cell lines, nestin co-localized incompletely with bundles of IFs containing other IF proteins (i.e. vimentin, glial filament, neurofilament). Nestin was ubiquitous in a wide variety of brain tumors, but was most prominent in gliomas. These studies document the existence of a human counterpart of rat nestin.

Distribution of the Nestin Gene Product in the Human Developing CNS

Cervical levels of six developing human spinal cords with gestational ages (GAs) of 6 to 40 weeks were immuno-histochemically stained with anti-nestin 129. The spinal cord of the 6 weeks GA human fetus was composed of three distinct layers, i.e. a layer of primitive neuro-epithelial cells lining the central canal, a mantle layer and an outer marginal layer. Anti-nestin 129 stained most cells in the primitive neuroepithelial layer, i.e. the presumptive multipotential stem cells that give rise to CNS neurons and glia. In addition, thin elongated radial glial fibers extending from the primitive neuro-

-22-

epithelial layer to the subpial region were stained by this antiserum. These neuroepithelial stem cells and radial glial fibers also expressed vimentin but not GFAP. Neuroblasts in the mantle layer (identified by their morphology and NF protein positively) did not express nestin.

At 11 weeks GA, cells in the ependymal layer were positively stained with anti-nestin 129, as were radial glial fibers. These radial glial fibers were also stained with the anti-vimentin MAb. Nestin immunoreactivity decreased at 17 weeks GA and disappeared at 20 weeks GA. From this gestational age to maturity, GFAP positive glial cells and NF positive neurons were seen in the spinal cord, but no elements other than endothelial cells were nestin positive.

The telencephalic germinal matrix of two cases with GAs of 17 and 20 weeks were examined with anti-nestin 129. At both time points, many presumptive neuroepithelial stem cells in the germinal matrix cells were nestin positive. These neuroepithelial precursor cells also were labeled by the anti-vimentin MAb, but no germinal matrix cells were stained by the MAbs to GFAP or NF proteins. At 40 weeks GA, a well defined germinal matrix was not evident, but clusters of morphologically immature cells (presumably residual, multipotential precursor cells) were seen in the subependymal region. Nestin immunoreactivity was recognized in only a few of these cells.

To determine if neuroepithelial precursors in the cerebellum also expressed nestin, four developing human cerebelli at GAs of 17 to 40 weeks were studied. Late in



-23-

development, the cerebellar cortex has four distinct layers, i.e. the internal and external granular layers, the Purkinje cell layer and the molecular layer. However, these layers were not yet evident at 17 weeks GA and Purkinje cells were not recognizable by morphological criteria. Nevertheless, immature NF positive cells were observed at 17 weeks GA in the superficial internal granular layer, and these cells probably correspond to nascent Purkinje cells. Nestin immunoreactivity was detected at this time in radial glial cells in the internal granular layer and in radial glial processes that extended to the external granular layer. These radial glial cells, which mature into the Bergmann glia of the adult cerebellum, also were stained by the anti-vimentin MAb (data not shown), but not by the anti-GFAP MAb. However, the anti-GFAP MAb did stain ependymal cells lining the fourth ventricle. At 20 weeks GA, nestin immunoreactivity in radial glia was markedly diminished and by 40 weeks GA, it was weakly present in only a few radial glia fibers, while the blood vessel endothelial cells at 40 weeks GA were clearly nestin positive. In contrast, radial glial fibers expressed both GFAP and vimentin (data not shown) at these time points.

## 25 Nestin Expression in Brain Tumor-derived Cell Lines

Indirect immunofluorescence studies of 9 well characterized human brain tumor derived cell lines showed variable nestin immunoreactivity in 6 or 7 primitive neuroectodermal tumor (PNET) cell lines, and very intense nestin staining in 2 glioma cell lines. Table 1

-24-

summarizes the data on the expression of nestin and other IF proteins in these cell lines.

TABLE 1

05

IMMUNOFLUORESCENCE ASSAY OF BRAIN TUMOR-DERIVED CELL LINES						
CELL LINES	NESTIN	VIMENTIN	GFAP	NF-L	NF-M	NF-H
Daoy	-	+	-	-	-	-
D283 Med	+	+	-	+	+	+
D341 Med	+	+	-	-	+	+
D384 Med	+	+	-	+	+	+
D425 Med	+	+	-	+	+	+
D458 Med	+	+	-	+	+	+
CHOP707m	+	+	-	+	-	-
U251 MG	+	+	+	-	-	-

15 Summary of the data obtained from the indirect immunofluorescence studies of each cell line. The patterns of staining are described as - = no staining, + = positive staining. NF-L, NF-M and NF-H are the low, middle and high molecular weight neurofilament subunits, respectively. The following

20 antibodies were used to obtain these data; nestin (rabbit anti-nestin antiserum 129), vimentin (V9), GFAP (2.2B10), NF-L (NR-4, RMS12), NF-M (RMd020, RM0254, H014), NF-H (DP1, TA51, RM024).

25 Except for CHOP707m, which was derived from a cerebral neuroblastoma (Baker, D.L. et al., Am. Neurol. 28:136 (1990)), the other 6 PNET cell lines were obtained from cerebellar medulloblastomas (see Hockfield, S. et

-25-

al., J. Neurosci 5:3310 (1985)) and citations therein). Daoy does not exhibit any evidence of glial or neuronal differentiation and it was nestin negative, while the other 6 PNET cell lines resembled that of embryonic neuroblasts (He, X. et al., J. Neuropathol. Exp. Neurol. 48:48 (1989)). Double immunofluorescent staining of D283 Med, which is the most differentiated PNET cell line, showed incomplete co-localization of nestin and NF proteins in bundles of IFs in the same cells. Similar observations were made using nestin and vimentin antibodies. CHOP707m expressed extensive nestin immunoreactivity and double immunofluorescence of nestin and vimentin in this cell line co-localized both proteins in the same cells, but the nestin positive filament bundles were located mainly in the perinuclear area while vimentin positive filament bundles were present throughout these cells. The Daoy line expresses vimentin, but not NF proteins, GFAP or other molecular markers of neurons or glia (He, X. et al., J. Neuropathol. Exp. Neurol. 48:48 (1989)). Hence, Daoy is the least differentiated PNET cell line, and it did not express nestin. The foregoing co-localization studies also indicated that the anti-nestin 129 did not cross react with vimentin or NF proteins.

Since nestin is expressed primarily in CNS stem cells and is eliminated during the progressive maturation of the progeny of these stem cells, the Daoy cell line may resemble a CNS precursor that subsequently develops into a vimentin and nestin positive CNS stem cell like those observed here in the human spinal cord at a GA of 6 weeks. Alternatively, tumor cell lines contain genetic

-26-

mutations and their phenotype may not fully replicate that of normal cells. However, the 2 glioma cell lines (U251 MG and U373 MG) resembled immature spinal cord radial glia (i.e. at 11 weeks GA) since these cells  
05 co-expressed nestin, GFAP and vimentin in the triple fluorescence studies.

Finally, Western blot studies were performed which showed that anti-nestin 129 labeled a single band in cytoskeletal extracts of U251 MG cells, and this immuno-  
10 band had a  $M_r$  similar to that of the bacterial TrpE-nestin fusion protein and lower than that of rat nestin expressed in P6 rat cerebellum. The significance of the difference in the  $M_r$  of rat and human nestin is unclear and will require further studies. Since these extracts  
15 contained vimentin, GFAP and other cytoskeletal proteins, these data further demonstrate the specificity of the anti-nestin antiserum for human nestin.

#### THE EXPRESSION OF NESTIN IN BRAIN TUMOR TISSUE

##### Brain Tumors Contain Nestin Positive Cells

20 ST15A cells express properties expected of the precursor to brain tumor cells as described in Example 3. Therefore, brain tumor cells were tested to determine whether they express features in common with ST15A, such as expression of nestin. As described in detail in  
25 Example 6, tissue from five medulloblastomas was obtained. Three of these samples came from recurrent tumors occurring in a 16 year old male, a 26 year old male and a 5 year old female. The remaining tumors were

-27-

newly diagnosed in an 11 year old male and a 10 year old female. The primary tumor in the 11 year old male was also used as a source of cultured cells.

05 Immunohistochemistry on sections of tissue from each  
of the five tumors studied showed regions with large  
numbers of nestin immunoreactive cells. The expression  
of nestin in medulloblastoma tissue was not uniform and  
in some cases parts of the tumor tissue showed a few  
nestin-positive cells among nestin-negative cells. The  
10 pre-immune serum showed no reaction with either human  
embryonic or tumor tissue. The nestin positive cells  
were small with a cytoplasmic distribution of the  
antigen. In one tumor, there were groups of aligned  
nestin positive cells which were very similar in their  
15 morphology to the embryonic neuroepithelium.

A cell line which was generated by growing cells out  
of the medulloblastoma from the 11 year old male was also  
tested for expression of nestin. Immunocytochemistry  
with polyclonal anti-nestin antiserum shows that these  
20 cells are uniformly positive for nestin expression and  
reveals the filamentous pattern characteristic of this  
intermediate filament protein. Western blotting using  
anti-nestin serum shows that the antigen detected in  
these cells comigrates with rat nestin.

25 Sections of normal fetal and adult human cerebellum  
were also stained with polyclonal antiserum against  
nestin. The fetal tissue showed many positive cells in  
the proliferative, external granular layer of the  
developing cerebellum and in the outer part of the  
30 internal granular layer. In contrast, there was no

-28-

immunoreactivity in cerebellum from a 2.5 year old. These results are consistent with the stem cell specific expression of nestin seen in developing mouse and rat brain.

#### 05 Utility

The adult cell types of the brain express distinct intermediate filament proteins: neurofilaments are expressed by neurons, peripherin by a subset of neurons (Leonard, Gorham, et al., (1988)), and glial fibrillary acidic protein by astrocytes. The expression of nestin defines an earlier stage in the pathway of intermediate filament gene expression. Early embryonic cells are nestin negative but express certain cytokeratins (Jackson, B.W., C. Grund et al., Differentiation 20: 203-216 (1980)). During neurulation neurectodermal cells become nestin positive. Upon terminal differentiation, CNS cells stop producing nestin and express class III or IV intermediate filaments typical of their differentiated cell type.

20 Nestin positive neurectodermal cells have been shown to be capable of differentiating into neuronal, glial and muscle cells (i.e., the different cell types found in tumors). Nestin expression has also been directly detected in tumor tissue and in a medulloblastoma derived cell line. Nestin positive cells derived from tumors are likely to be renewing stem cells which give rise to the differentiated cells in the tumor.

25 The detection of nestin expression in brain cells obtained from an adult using known methods is indicative of the presence of a brain tumor or of neural precursor

30

-29-

cells capable of developing into a brain tumor. For the purposes of the subject invention, an "adult" is an individual whose central nervous system is developed and, therefore, contains differentiated cells. For example, 05 the central nervous system of human neonates is in continual development up until the age of about one year. Therefore, a human adult, for the purposes of the subject invention, is an individual who is older than about one year.

10 Nestin mRNA expression in a brain tissue sample can be detected using a DNA probe; this is indicative of expression of nestin, which is, in turn an indication of a predisposition to the development of a brain tumor or of the presence of a brain tumor in the individual. For 15 example, a sample of brain tissue from of an individual can be hybridized to a DNA probe which is complementary to all or a portion of the nestin gene. Detection of hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain 20 tumor.

Alternatively, expression of the nestin protein can be detected using polyclonal antibodies, (e.g., anti-nestin antiserum 129) or monoclonal antibodies (e.g., Rat 401). For example, cerebral spinal fluid or a serum 25 sample from the brain of an individual can be stained with anti-nestin antibodies. Detection of stained cells is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor.

The finding that neural precursors to tumors express 30 nestin also has important therapeutic implications. For example, the nestin marker enables further investigation

-30-

into the proliferation and differentiation of the stem cells of a brain tumor, so that existing chemotherapies can be implemented at an earlier stage in the tumor development and so that new methods of inhibiting tumor development can be developed.

The present invention will now be illustrated by the following examples, which are not to be seen as limiting in any way.

#### EXEMPLIFICATION

##### 10 EXAMPLE 1 Cloning and Characterization of the Rat Nestin Gene

###### Construction and Screening of Libraries

15  $\lambda$ gt10 and  $\lambda$ gt11 cDNA libraries were constructed from oligo-dT-primed poly(A)+ RNA from embryonic day 15 dissected rat CNS according to procedures which were slightly modified from those described (Young, R.A. and R.W.Davis, Proc. Natl. Acad. Sci. USA. 80: 1194-1198 (1983)). The second strand cDNA was synthesized using the RNase H modification of the Okayama-Berg method (Gubler, U. and R. Hoffman, Gene 25: 263-269 (1983)). After induced protein expression, the  $\lambda$ gt11 library was screened with the monoclonal antibody Rat 401 (undiluted hybridoma supernatant) and positive clones were visualized with an anti-mouse second antibody conjugated to alkaline phosphatase (Promega Protoblot). Immunopositive clones were selected and the cDNA insert from the clone giving the strongest hybridization signal,  $\lambda$ gt11.401:16,



-31-

was <sup>32</sup>P-labelled (Feinberg, A.P. and B. Vogelstein, Anal. Biochem. **137**: 266-267 (1983) and used as probe to screen the  $\lambda$  gt10 library (Benton W.D. and R.W. Davis, Science. **196**: 180-182 (1977); Thomas, P.S. Proc. Natl. Acad. Sci. USA. **77**: 5201-5205 (1980)). Several clones were identified. A commercial rat genomic DNA library (Clontech) in the vector Charon 4A was screened with <sup>32</sup>P-labelled nestin cDNA inserts as above. A number of hybridizing clones were obtained, and a genomic map comprising 20 kb of the nestin gene region was established by restriction mapping of the genomic clones, cross-hybridizations, and sequencing of selected genomic subclones.

#### DNA Sequencing and Computer Analysis

cDNA inserts and genomic subclones were sequenced after subcloning into M13mp18, M13mp19, or Bluescript KS+ (Stratagene). M13 phage particles and Bluescript were grown and circular DNA prepared according to standard procedures. Both single stranded (M13) and double stranded (Bluescript) sequencing was performed using a modified T7 DNA polymerase (Sequenase, USB), as suggested by the manufacturer. Approximately 90% of the gene, including all exons and cDNA inserts, was sequenced on both strands. Ambiguous regions were further resolved using deoxyinosine in the standard Sequenase protocol. Computer analysis was performed using the University of Wisconsin Genetics Computer Group program package, including the FASTP and TFASTA algorithms (Devereux, Haeberli et al., 1984)). The complete sequence, including introns, will be stored in the EMBL Database.

-32-

### Isolation of RNA and Northern Blotting

Tissue was rapidly dissected and quickly frozen in liquid nitrogen. After homogenization in a guanidinium thiocyanate containing buffer, total RNA was isolated by centrifugation through a CsCl cushion (Chirgwin, Przybyla *et al.*, 1979)). Poly(A)+ RNA was isolated by an oligo-dT chromatography column (Aviv and Leder, 1972) or by a poly(A)+ RNA extraction kit (Fast Trak, Invitrogen). For Northern blot analysis RNA was denatured and electrophoresed in 1% agarose gels containing 2.2 M formaldehyde in 1X MOPS buffer (Maniatis, Fritsch *et al.*, 1982) at +4°C for 8 hours. The RNA was then transferred to nitrocellulose filters (GeneScreen Plus, New England Nuclear) (Thomas, 1980)). The filters were prehybridized and hybridized (50% formamide, 0.75M NaCl, 100 mM Tris pH 7.8, 5 mM EDTA, 1% sodium dodecyl sulphate at 42°C for 16 hours) with  $2 \times 10^6$  cpm/ml of the  $^{32}$ P-labeled nestin cDNA insert from  $\lambda$  gt10.401:16 (specific activity  $3 \times 10^8$  cpm/ug). After hybridization, the filters were washed (final wash - 2 x 30 minutes at 65°C in 30 mM NaCl, 4 mM Tris pH 7.8, 0.2 mM EDTA and 0.25% SDS) and exposed to x-ray film with intensifying screens. Quantity and integrity of the RNA were monitored by rehybridizing the blots with a cDNA probe from the human beta-tubulin gene under identical conditions. The size of the hybridizing RNA was determined by running RNA molecules of known size (HMW RNA ladder, BRL) in parallel.

### S1 Nuclease Mapping and Primer Extension

To generate a probe for S1 nuclease protection experiments, we amplified a region of a genomic clone

-33-

between two synthetic primers using the polymerase chain reaction (kit from Perkin-Elmer Cetus). The resulting fragment was 315 bp long and located between bp -121 and +194 (bp +1 indicates the subsequently defined initiation of transcription). After  $^{32}\text{P}$  end-labelling, 300 fmole of the fragment was hybridized to 50 ug of E15 CNS rat total RNA or 50 ug of yeast total RNA according to standard procedures, and the S1 resistant hybrids separated on denaturing polyacrylamide gels (Maniatis, T., E.F. Fritsch et al., Molecular cloning: A laboratory manual (Cold Spring Harbor, New York: (1982)).  $^{32}\text{P}$  end-labeled restriction digested plasmids and sequencing reactions of known DNAs were used as size standards. The dried gels were exposed to x-ray film. Primer extensions were performed by hybridizing 125 fmole of one of the oligonucleotides used in the polymerase chain reaction (pos +170 - +194) to 10 ug of poly(A)+ RNA from rat E15 CNS according to standard procedures. The reaction product was subjected to gel electrophoresis as in the S1 nuclease experiments.

#### Immunocytochemistry

For all experiments cells from the immortalized cell line ST15A were grown for 2 days (to 40% confluence) in Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum on sterile, polyornithine coated glass cover slips, as described in U.S. Patent Applications Serial Nos. 07/603,803; 07/201,762 and 180,548. All antibody incubations lasted 60 minutes at room temperature and were followed by three rinses in PBS (0.14M)

-34-

NaCl, 2.7 mM KCl, 8 mM  $\text{Na}_2\text{HPO}_4$  and 8 mM  $\text{KH}_2\text{PO}_4$  pH 7.2). In the double-label experiments with nestin and tubulin antibodies the ST15A cells were briefly rinsed in PBS, presoaked in microtubule stabilizing buffer (MTSB - 2M glycerol, 0.1 M PIPES pH 6.9, 1 mM  $\text{MgSO}_4$ , 2 mM EDTA) for 2 minutes and then fixed for 5 minutes in 4% para-formaldehyde, 0.5% NP-40 in MTSB for 5 minutes. After two brief rinses in PBS and 5 minutes incubation in blocking solution (0.75% Triton X-100, 500 ug/ml bovine serum albumin, 1% horse serum and 0.05% sodium azide), the cells were incubated with a polyclonal rabbit-anti-tubulin antiserum (kind gift of Dr. Frank Solomon) diluted 1:10000 in PBS. This was followed by an incubation with a fluorescein-conjugated goat-anti-rabbit IgG antiserum (Cappel, No. 1612-0081) diluted 1:100 in PBS. The cells were then postfixed in 95% ethanol/5% acetic acid for 5 seconds and rinsed three times in PBS. This was followed by an incubation with undiluted hybridoma supernatant from the monoclonal mouse-anti-nestin antibody (Rat 401). Finally, the cells were incubated with a rhodamine-conjugated polyclonal goat-anti-mouse IgG antibody (1:100 dilution, Cappel No. 2611-0231). After two brief rinses in water, the cells were covered with Immumount (Shandon) and inverted onto clean glass microscope slides. As control experiments for potential cross-hybridization and spillover between the fluorescein and rhodamine spectra in the fluorescence microscope, the first antibodies (rabbit-anti-tubulin and mouse-anti-nestin) were omitted in two separate experiments under otherwise identical conditions. No cross-hybridization

-35-

or significant spillover was observed. In the colchicine experiments the ST15A cells were grown as above and treated with colchicine at 25 ug/ml for 24 hours followed by fixation and staining as described here.

05 In the actin/nestin double-label experiments ST15A cells were grown as above and fixed in 4% paraformaldehyde, 0.1% Triton X-100 in PBS for 5 minutes. After blocking as described above, the cells were incubated with 0.16 uM rhodamine-conjugated phalloidin, as  
10 suggested by the manufacturer (Molecular Probes). Rinses, postfixation and staining with the nestin monoclonal antibody were performed as described above, with the exception that a fluorescein-conjugated sheep-anti-mouse IgG second antibody (diluted 1:100, Cappel No.  
15 1606-3152) was used. No cross-hybridization or spillover was observed after omitting the rhodamin-conjugated phalloidin or the nestin antibody. In the cytochalasin B experiments the ST15A cells were treated with cytochalasin B at a final concentration of 10 ug/ml for 60  
20 minutes, otherwise all immunostaining procedures were identical.

In the vimentin/nestin double-label experiments ST15A cells were fixed in 4% paraformaldehyde, 0.1% Triton X-100 in PBS for 5 minutes. After blocking, the  
25 cells were incubated with a polyclonal goat-anti-vimentin antiserum (dilution 1:20, MILES 65-794) followed by a fluorescein-conjugated rabbit-anti-goat IgG second antibody (dilution 1:100, ICN 65-176). Incubation times and rinsing procedures were as described above. Post-  
30 fixation and incubation with the anti-nestin antibody

-36-

were performed as in the tubulin/nestin experiments. No cross-hybridization or spillover was detected after omitting the anti-vimentin and anti-nestin antibodies, respectively. In all experiments the mounted cells were  
05 photographed on Kodak Ectachrome 400 ASA using a 63X objective on a Zeiss fluorescence microscope.

EXAMPLE 2 Production of the Rat 401 Monoclonal Antibody

Developing monoclonal antibodies were generated to fixed spinal cord from embryonic day 15 (E15) rats.  
10 Timed pregnant female rats were obtained from Taconic Animal Supply Co. and housed until the appropriate gestation age. Uteri were dissected from pregnant animals into ice-cold phosphate buffer (pH 7.4), individual embryos were removed and the spinal cord was  
15 dissected free of other tissue into fixative (4% paraformaldehyde in 0.1 M phosphate buffer).

The immunization and fusion protocol has been described previously (McKay and Hockfield, Proc. Natl. Acad. Sci. USA 79: 6747-6751 (1982)). Briefly, BALB/c  
20 mice received two intraperitoneal immunizations with fixed tissue homogenized in saline and suspended in an equal volume of Freud's complete adjuvant and a final intravenous boost of unfixed tissue with adjuvant. Spleen cells from immunized mice were fused with NS1  
25 myeloma cells and resulting hybrid cell lines were screened immunohistochemically on 100  $\mu$ m-thick Vibratome sections of 4% paraformaldehyde-fixed E15 rats. Lines producing antibodies of interest were cloned in soft agar or by limiting dilution and antibody subclass was deter-  
30 mined by the Ouchterlony method (Ouchterlony and Nilsson,

-37-

Immunodiffusion and Immunelectrophoresis. In Handbook of Experimental Immunology D.M. Weir, ed., Blackwell Scientific Publications, Oxford (1978).

#### Immunohistochemistry

05        Tissue for immunohistochemistry was fixed either by  
intravascular perfusion or by immersion. All postnatal  
material was fixed by intravascular perfusion (under  
ether anesthesia) with 4% paraformaldehyde at pH 7.4 for  
light microscopy or with either 4% paraformaldehyde at pH  
10 10.0 or 4% paraformaldehyde, 0.1% glutaraldehyde at pH  
7.4 for electron microscopy. The CNS was dissected out  
and stored in 0.1 M phosphate buffer (pH 7.4) with 0.1%  
sodium azide. For prenatal material, pregnant females  
were ether anesthetized, uteri were dissected into cold  
15 0.1 M phosphate buffer and the females were killed by  
cervical dislocation. Embryos were dissected indi-  
vidually into Sylgard-coated Petri dishes and pinned in  
place. Intravascular perfusion was performed by opening  
the skin of the thoracic cavity, nicking the right atrium  
20 with a fine forceps and placing a 25 gauge needle into  
the left ventricle. One to 5 ml of fixative (as above)  
were injected into the heart. The embryos were placed in  
a vial containing the same fixative for 4 hr and then  
stored in phosphate buffer. E12 to term embryos were  
25 fixed by perfusion (using a 30 gauge needle for E12  
embryos); E10 and E11 embryos were fixed by immersion in  
fixative. E13 and older animals were sectioned at 50 to  
100  $\mu$ m on a Vibratome without additional support. E10 to  
E12 embryos were embedded in 15% gelatin before  
30 sectioning.

-38-

Immunohistochemical reactions were performed on free-floating sections by sequential incubations in monoclonal antibody as full-strength supernatant (12 to 20 hr); horseradish peroxidase (HRP)-conjugated goat anti-mouse antibody at a dilution of 1:100 in tissue culture medium with 10% serum; and 3,3'-diaminobenzidine (DAB; 0.025%) with  $H_2O_2$  (0.002%). Peroxidase-conjugated lectins (Sigma Chemical Co.) were used at serial dilutions in the same manner with the omission of the peroxidase-conjugated goat anti-mouse antibody. For light microscopy, 2% Triton X-100 was included in both antibody solutions. For electron microscopy sections carried through this procedure (without Triton X-100) were post-fixed in 2%  $O_2O_4$  embedded in Epon/Araldite between plastic coverslips, and cut at 1  $\mu m$  and 0.12  $\mu m$  for correlative light and electron microscopy. For high resolution light microscopy, sections (Triton-treated) were embedded in plastic as for electron microscopy and sectioned at 2  $\mu m$ .

## 20 Immunoblots

The apparent molecular weights of antigens recognized by antibodies were determined using the method of Towbin *et al.*, Proc. Natl. Acad. Sci. USA 76: 4350-4354 (1979)). Proteins were extracted from unfixed tissue with sodium dodecyl sulfate (SDS) and  $\beta$ -mercaptoethanol and run on an SDS-polyacrylamide gel. The proteins were electroblotted onto nitrocellulose paper which was then exposed to primary (1 hr) and secondary (1 hr) antibodies. HRP was visualized with 4-chloronaphthol.



-39-

EXAMPLE 3 Characterization of Rat Nestin from the  
ST15A Cell Line

ST15A cells were derived from rat postnatal day 2 cerebellum after infection of a primary culture with a  
05 retrovirus transducing the temperature sensitive, tsA58 allele of SV40 T antigen (Frederiksen et al., 1988)). ST15A cells were plated on tissue culture plastic coated with polyornithine and passaged at the permissive temperature, 33°C, in Dulbecco's modified Eagle's medium (DME)  
10 supplemented with 2 mM L-glutamine, 1 mM sodium pyruvate and 15 mM HEPES, pH 7.2, containing 10% fetal calf serum (FCS). For the experiment in Figure 8, confluent cells (labelled 'day 0') were transferred into DME/10% horse serum (HS) and cultured at 33°C or 39°C as marked for the  
15 indicated time. 39°C is the non-permissive temperature for tsA58.

Primary cultures were prepared from P4 rat cerebellum as previously described (McKay, R.D.G. et al.,  
20 Cold Spring Harbor Symp. Quant. Biol. (1990). Briefly, tissue was dissociated by trypsinization followed by trituration. Cells were plated on polyornithine-coated coverslips in DME/10% FCS/10% HS and grown at 39°C for 24 hours. The medium was changed to DME/10% HS and culture continued for 6 days.

25 The medulloblastoma cell line was derived by growing cells from dispersed, fresh medulloblastoma tissue. The tissue was removed from a previously untreated tumor in a 10 year old female. The cell line has not been cloned. Cells were passaged in Ham's F12 supplemented with 20mm  
30 L-glutamine and 10% FCS.

-40-

Immunocytochemistry

Cells for immunocytochemistry were fixed in 4% paraformaldehyde buffered with 50 mM sodium phosphate, pH 7.2, for troponin staining or buffered with 50mM sodium borate, pH 9.5, for nestin staining. The antibodies used were mouse monoclonal Rat 401 (Hockfield and McKay, supra (1985)), mouse monoclonal antibody against troponin T (Sigma T-6277), and rabbit polyclonal serum raised against the C-terminal region of nestin expressed in bacteria. Fixed cells were incubated in primary antibody for one hour, rinsed 3 times in PBS, incubated for one hour with flourescein-conjugated goat anti-rabbit serum (Organon-Tetnika) and mounted in Immumount (Shandon, Inc.) with 2% DABCO. Medulloblastoma tissue and normal human cerebellum were frozen and subsequently fixed in formalin. 50 um sections were cut and stained with rabbit polyclonal serum raised against the C-terminal region of nestin.

Isolation of RNA and Northern blotting

Cultured cells were scraped into 5M guanidine thiocyanate/25 mM citrate, pH 6.8/ 0.5% N-lauryl sarcosine/0.1  $\beta$ -mercaptoethanol and RNA pelleted through a CsCl step gradient (Chirgwin et al., 1979). 20 ug of total RNA was electrophoresed through a 1% agarose/2.2M formaldehyde gel and transferred to nitrocellulose. Prehybridization and hybridization were in 50% formamide/5xSSC/5X Denhardt's/0.1% SDS/0.1 mg/ml salmon sperm DNA at 42°C. Filters were washed in 2xSSC/0.1% SDS at room temperature and in 0.1xSSC/0.1% SDS at 55°C and

-41-

exposed to X-ray film with an intensifying screen. Filters were stripped by washing 6 times for 3 minutes each in boiling 0.05xSSC/0.1% SDS/20 mM EDTA, pH8, and rehybridized. Probes were prepared for hybridization by random priming on fragments isolated on low melting temperature agarose gels (Feinberg and Vogelstein, supra (1984)). The following plasmids were used to generate probes: mouse myoD1 cDNA pEMC11s (Davis, R.L. et al., Cell. 51: 987-1000 (1987); mouse myogenin cDNA p65 (Wright et al., supra (1989); rat nestin cDNA p401-16; human B-actin cDNA (Gunning, P. et al., Mol. Cell. Biol. 3: 787-795 (1983)).

#### Western Blotting

Proteins were extracted from cultured cells or rat postnatal day 5 limb muscle (Sprague-Dawley; Taconic, Inc.) in the presence of aprotinin and PMSF in 2% SDS, 5 mM EDTA and 15% glycerol in 0.1 M Tris, pH 6.8. 50 ug of total protein was run in each lane of an SDS reducing gel according to Laemmli (Laemmli 1970); proteins were transferred to Immobilon P membranes (Millipore) using a Hoeffer transfer apparatus at 50 volts for 4 hours. Filters were blocked in 5% BLOTTO (Johnson, D.A. et al., Gene Anal. Tech. 1: 3-8 (1984)) overnight. Rat 401 hybridoma supernatant was used undiluted; troponin T antibody was diluted 1:500; polyclonal anti-nestin antiserum was diluted 1:1000; HRP conjugated secondary antibodies were purchased from BioRad.

-42-

**EXAMPLE 4 Characterization of Human Nestin**

To characterize human nestin, a new antiserum, designated anti-nestin 129, was raised to recombinant rat nestin expressed in E. coli since the rat-401 mAb failed to recognize human nestin in immunohistochemical and immunochemical assays.

**Production of Anti-nestin Antiserum**

The insert from the clone  $\lambda$ gt10 401:16 (Lendahl, U. et al., Cell 60: 585 (1990)) was isolated and ligated into the pATH1 vector (Tzagaloff, A. et al., J. Biol. Chem. 262: 17163 (1986)), producing a clone in which the bacterial TrpE protein (37.2 kD) was fused at its C terminus with the last 1197 amino acids of nestin. The fusion protein was induced in E. coli strain HB101 with 20  $\mu$ g/ml indoleacrylic acid (Sigma I1625) for 4 hours as described in (Tapscott, S. et al., Science 242: 405 (1988)). The TrpE-nestin fusion protein migrates at greater than 200 kD in 6% SDS PAGE gels, making it the largest protein in the bacterial lysate. The fusion protein was purified by cutting the top band from the gel. Anti-nestin antiserum 129 was produced by injecting the fusion protein subcutaneously into a female NZW rabbit. The characterization of anti-nestin 129 and its specificity for human nestin is described here.

First, this antiserum was compared to the Rat-401 MAb in immunochemical and immunohistochemical studies of postnatal day 6 (P6) rat cerebellum. Then, anti-nestin 129 was used to probe human CNS tissues, tumors and tumor derived cell lines. To determine if nestin co-localized with other IF proteins, double and triple immuno-

-43-

fluorescence experiments also were performed using anti-nestin 129 and antibodies to vimentin, glial fibrillary acidic protein (GFAP) and neurofilament (NF) triplet proteins. Finally, Western blots were performed  
05 on cytoskeletal extracts of some of the tumor derived cell lines to identify human nestin and to compare it with rat nestin as well as with the class III and IV IF proteins expressed by CNS cells.

Specificity of Anti-Nestin 129 in the Rat

10 The rabbit anti-nestin antiserum produced as described herein, i.e. anti-nestin 129, yielded results identical to those produced by the Rat-401 MAb when both were used to probe P6 and adult rat CNS tissues by immunoblots and immunohistochemistry. On immunoblots of  
15 cytoskeletal extracts from P6 rat cerebellum separated by SDS-PAGE, anti-nestin 129 identified a band with an apparent  $M_r$  of 240 kD. This band was identical to the band labeled by Rat-401. Further, both antibodies labeled similar bands in lysates of the induced bacteria  
20 producing the TrpE-nestin fusion protein.

In immunohistochemical studies of the P6 rat cerebellum, anti-nestin 129 labelled radial glial cells very intensively. The processes of these cells radiated from the internal granular layer to the external granular  
25 layer. These radial glia also were labeled with Rat-401 and the anti-vimentin MAb. Neither of the anti-nestin antibodies stained immature Purkinje cells or other neurons. In contrast to the P6 rat cerebellum, the anti-nestin 129 and Rat-401 antibodies only stained blood  
30 vessel endothelial cells in the adult rat cerebellum,

-44-

while the anti-vimentin MAb (V9) labelled blood vessels, white matter astrocytes and radial glial fibers. Furthermore, nestin immunoreactivity was demonstrated by both anti-nestin antibodies in immature skeletal muscle of the P6 rat as well as in Schwann cells of adult rats (data not shown).

These studies demonstrate that this new anti-nestin antiserum yields immunohistochemical results in the human developing spinal cord, cerebrum and cerebellum that were nearly identical to those produced by Rat-401 and the anti-nestin antiserum in the rat.

#### EXAMPLE 5 Cloning of the Human Nestin Gene

The human nestin gene was isolated using low-stringency DNA hybridization of a human genomic bacteriophage lambda library with a rat nestin probe.

A genomic library in the vector EMBL 3 made from partially Sau3A-digested human genomic DNA (Clontech) was screened using a  $^{32}\text{P}$ -labelled probe derived from the rat nestin gene starting at 202 nucleotides upstream of the translation start and ending at nucleotide 483 after the translation start (Lendahl, et al., 1990) under the following conditions: prehybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 micrograms/ml salmon sperm DNA at 65°C for 1 hour; hybridization in the same conditions except for using 50 micrograms/ml salmon sperm DNA and hybridizing for 16 hours. The filters were then washed for 2 X 30 minutes at 48°C in 2X SSC and 0.25% SDS.

One positive clone out of  $1.5 \times 10^6$  plaques screened contained the entire human nestin gene and was

-45-

chosen for further analysis. This clone was purified and grown in large scale (Maniatis, et al., 1989). A restriction map was established and EcoRI fragments were subcloned in the plasmid vector Bluescript KS I (Stratagene).

05

**EXAMPLE 6 Nestin Expression in Brain Tumor Samples**

Immunohistochemical data on the expression of nestin, other IF proteins and synaptophysin (SYP) by a diverse group of 34 CNS tumors are summarized in Table 2.

**TABLE 2 Immunoperoxidase staining of brain tumors**

No.	Age(yr)	Sex	Material	Location	Nestin	Vim	GFAP	NP	SYP
<b>PNETs</b>									
<b>PNET+N/G</b>									
1	1	M	F	posterior fossa	+	+++	+	+	-
2	1	M	F	posterior fossa	+	+++	+	+++	+++
3	5	F	F	posterior fossa	++	+	+	+	++
4	-1	M	F	posterior fossa	++	++	+	+	+++
5	1 month	F	F	posterior fossa	++	++	+	+++	+
6	5	M	F	posterior fossa	++	++++	++	+	+
7	8 months	F	F	posterior fossa	+++	+++	++	+	+
8	1	F	F+P	posterior fossa <sup>d</sup>	++	++	+	+	+++
<b>PNET+N</b>									
9	13	F	F	posterior fossa <sup>b</sup>	-	++++	-	++	++
10	6	M	F	posterior fossa	-	+	E	++	++
<b>PNET+G</b>									
11	11	M	F	posterior fossa	+	+++	++	P	+
12	4	F	F	posterior fossa	+++	++++	++	P	+++
<b>PNET+NOS</b>									
13	2	M	F	pineal	++	++	-	-	++
14	11	M	F	posterior fossa <sup>b</sup>	-	+	-	-	++
15	14	M	F	posterior fossa <sup>b</sup>	+	+++	-	-	+
<b>Astrocytic tumors</b>									
16	2	F	F	posterior fossa <sup>c</sup>	++	++++	++++	-	-
17	6	F	F+P	posterior fossa <sup>c</sup>	-	++++	+++	-	-
18	6 months	M	F	hypothalamus <sup>c</sup>	++	+++	+++	-	-
19	17	M	F+P	posterior fossa <sup>d</sup>	-	+++	+++	-	-
20	3	M	F	frontal <sup>e</sup>	+++	NA	+	-	NA
21	6	M	F+P	occipital <sup>f</sup>	+++	+++	+++	-	-
22	13	M	F	posterior fossa <sup>f</sup>	+++	+++	+++	-	-
<b>Ependymomas</b>									
23	3	F	F	parietal	++	++++	+	-	-
24	2	M	F	posterior fossa	+	-	+	-	-
25	3	M	F+P	posterior fossa	++	++++	+++	-	-
26	5	M	F+P	posterior fossa	++	++++	++	-	-
27	15	F	F+P	cerebral hemisphere <sup>g</sup>	++	++++	+	-	-
28	8	F	F+P	parietal <sup>h</sup>	+++	+++	++	-	-
<b>Choroid plexus papillomas</b>									
29	1	M	F	posterior fossa	-	+++	+	-	-
30	2	M	F	lateral ventricle <sup>i</sup>	-	+++	-	-	-
<b>Gangliogliomas</b>									
31	2	F	F	temporal	++	+++	+++	+	-
32	1	F	F	occipital	++	++	+++	+	-
<b>Meningiomas</b>									
33	8	M	F	posterior fossa	++	++	-	-	-
34	27	M	F+P	anterior fossa	+	+++	-	-	-



-47-

Legend to Table 2

Summary of the data obtained from the immunoperoxidase studies of each of the brain tumor samples. Results from frozen (F) and paraffin (P) material are combined in the cases in which both kinds of material were studied. The patterns of staining are described as - = no staining, + = positive in less than 5% of tumor cells, ++ = positive in between 5 - 50% of tumor cells, +++ = positive in between 50 - 95% of tumor cells, ++++ = positive in more than 95% of tumor cells, E = positive cells equivocal, i.e. reactive astrocytes of tumor cells, P = interstitial filamentous staining in the neuropil. The following antibodies were used to obtain these data; nestin (rabbit anti-nestin antiserum 129), vimentin (V9), GFAP (2.2B10), NF (RMS12, RMd020, RM0254, H014, DP1, TA51, RM024). NA = not available due to limited material; PNET+N,G = PNET with neuronal and glial differentiation; PNET+N = PNET with neuronal differentiation; PNET + G = PNET with glial differentiation; PNET-NOS = PNET, not otherwise specified. <sup>a</sup>reccurent tumor of case 7, <sup>b</sup>autopsy case, <sup>c</sup>benighn astrocytoma <sup>d</sup>genighp astrocytoma + angioma, <sup>e</sup>anaplastic astrocytoma <sup>f</sup>glioblastoma multiforme, <sup>g</sup>mixed ependymo-astrocytoma, <sup>h</sup>anaplastic enpendymoma, <sup>i</sup>choroid plexus carcinoma.

- 25 The results from frozen and ethanol-fixed paraffin-embedded samples from the same biopsy are combined in the 10 cases from which these paired samples were available. Notably, the immunohistochemical results obtained from each of these paired samples were identical, although the
- 30 primary antibodies were used at slightly higher dilutions in the frozen material compared with the paraffin material.

- PNETs The anti-nestin antiserum stained tumor cells in 12 of the 15 PNETS examined here. Nestin immunore-
- 35 activity was found in the cytoplasm of individual neoplastic cells. Reactivity was also observed in the coarse processes of large stellate cells, as well as in

-48-

blood vessel endothelial cells. In the PNETs with an insular architecture, nestin reactive cells were found both within and outside the islands. Since PNETs express SYP and other neuroendocrine markers as well as all classes of IF proteins (Gould, V.E. et al., Human Pathol. 21:245 (1990)), the PNETs were probed for SYP. These studies demonstrated SYP in 14 of 15 PNETs. Vimentin was detected in all the PNETs. Ten of the 15 PNETs expressed GFAP and 10 expressed 1 or more NF proteins. Double immunofluorescence studies of nestin and NF proteins, as well as nestin and GFAP on frozen sections showed their co-localization in the same cells (data not shown). The immunohistochemical data on GFAP and NF proteins suggested that 15 PNETs could be classified into 4 subtypes, i.e. 3 cases of PNET-NOS (not otherwise specified), 2 cases of PNET+G (PNET with glial differentiation), 2 cases of PNET+N (PNET with neuronal differentiation) and 8 cases of PNET+G,N (PNET with glial and neuronal differentiation). All 8 cases of PNET+G,N showed nestin immunoreactivity and 2 cases of PNET+G expressed nestin. On the other hand, 2 cases of PNET+N were negative for nestin. These results may indicate that PNETs with neuronal differentiation tend to lack nestin expression in contrast to PNETs with glial differentiation. This phenomenon paralleled normal differentiation in that postmitotic neuroblasts did not express nestin whereas radial glial cells did, even at relatively late developmental stages. Two of 3 cases of PNET-NOS showed nestin positive tumor cells. This immunohistochemical phenotype might represent primitive neuro-epithelial cells or immature radial glial cells in the

-49-

developing CNS. The other PNET-NOS case was negative for nestin and it might correspond to extremely immature CNS precursor cells just like the most undifferentiated PNET cell line (i.e. Daoy).

05        Astrocytic tumors Five of 7 astrocytic tumors displayed positivity for nestin in the perinuclear cytoplasm of the tumor cells as well as in the cytoplasmic processes of these cells. Additionally blood vessel walls were nestin positive. Two of 4 benign  
10 astrocytomas were negative for nestin. One anaplastic astrocytoma and 2 glioblastomas showed strong nestin positivity. In the glioblastomas, multinuclear giant cells were stained strongly.

Ependymomas Six tumors were histologically  
15 diagnosed as ependymomas and all showed nestin positivity. Immunoreactivity for nestin was found in the tumor cells which formed perivascular pseudo-rosetts. One ependymoma (with marked nuclear pleomorphism, necrosis and a few mitoses) was diagnosed as an ana-  
20 plastic ependymoma and it demonstrated the strongest nestin immunoreactivity among the 6 ependymomas. Expression of nestin described here indicates that nestin is the dominant IF protein of early CNS precursor cells. However, nestin is rapidly extinguished in neurons, glia  
25 and other CNS cell types derived from these precursors. The mechanisms responsible for these rapid changes in IF protein expression are unknown.

         In contrast to the restricted expression of nestin in normal developing CNS cells, this IF protein was  
30 ubiquitously present in a wide variety of CNS neoplasms.

-50-

Nestin was expressed in a subset of PNETs, but was more abundant in gliomas. The presence of nestin in diverse types of neuroepithelial tumors is similar to vimentin, but distinct from the distribution of GFAP and NF

05 proteins in CNS tumors.

-51-

CLAIMS

1. Isolated DNA encoding a protein of mammalian origin whose expression distinguishes neural multipotential stem cells from neuronal, glial and muscle cells.
- 05 2. Isolated DNA of Claim 1 wherein the protein is nestin.
3. Isolated DNA of Claim 2 having the nucleotide sequence of all or a portion of the nucleotide sequence of Figure 1 or Figure 2.
- 10 4. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 1.
5. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 2.
- 15 6. Isolated DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 1.
7. Isolated DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 2.
- 20 8. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 1 which encodes a protein of mammalian origin whose expression distinguishes neural multipotential stem cells from neuronal, glial and muscle cells.
- 25

- 52 -

9. Isolated DNA having a nucleotide sequence corresponding to all or a portion of the nucleotide sequence of Figure 2 which encodes a protein of mammalian origin whose expression distinguishes  
05 - neural multipotential stem cells from neuronal, glial and muscle cells.
10. A DNA probe for the detection of a brain tumor, the probe comprising DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 1  
10 or to all or a portion of a DNA sequence sufficiently homologous to the DNA sequence of Figure 1 that it encodes a protein which is indicative of a predisposition to the development of a brain tumor in an individual.
- 15 11. A DNA probe for the detection of a brain tumor, the probe comprising DNA which hybridizes to all or a portion of the DNA sequence represented in Figure 2 or to all or a portion of a DNA sequence sufficiently homologous to the DNA sequence of Figure 2 that  
20 it encodes a protein which is indicative of a predisposition to the development of a brain tumor in an individual.

-53-

12. A method of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an individual, comprising:
- 05 a) obtaining a sample from the brain of an individual;
  - b) treating the sample in a manner that renders DNA present in the sample available for hybridization with a complementary DNA sequence thereby producing a treated sample;
  - 10 c) contacting the treated sample with at least one probe which is a DNA sequence which hybridizes with all or a portion of a DNA sequence represented in Figure 1 or Figure 2; and
  - 15 d) detecting the hybridization of DNA from the sample with the DNA probe; wherein hybridization is an indication of a predisposition to the development of a brain tumor or the presence of a brain tumor in the individual.
13. A method of diagnosing a predisposition to the development of a brain tumor or the presence of a brain tumor in an individual, comprising:
- 20 a) obtaining a sample from the brain of an individual;
  - 25 b) treating the sample in a manner that renders the cells available for staining with a nestin-specific antibody thereby producing a treated sample;
  - c) staining the cells of the treated sample with labelled anti-nestin antibodies; and
  - 30 d) detecting stained cells as an indication of a predisposition to the development of brain tumors or the presence of a brain tumor in an individual.

-54-

14. A method of Claim 13 wherein the antibodies are labelled with a label selected from the group consisting of fluorescein, rhodamine and peroxidase.
- 05 15. A method of Claim 13 wherein the nestin-specific antibody is a polyclonal antibody.
16. A method of Claim 15 wherein the polyclonal antibody is anti-nestin 129.
- 10 17. A kit for diagnosing a brain tumor, comprising at least one nucleic acid probe, which hybridizes with a DNA sequence represented in Figure 1 or Figure 2.
18. A kit for diagnosing a brain tumor, comprising a first antibody which is nestin-specific and a second antibody which binds to the first antibody and is labelled.
- 15 19. A nestin-specific antibody or a pharmaceutical composition comprising a nestin-specific antibody formulated in a solution for parenteral administration for use in in vivo diagnosis, e.g., tumor imaging.
- 20 20. Use of a nestin-specific antibody for the manufacture of a diagnostic agent for in vivo tumor imaging.



Figure 1

1  
22  
30  
38  
46  
54  
62  
70  
78  
86  
94  
102  
110  
118  
126  
134  
142  
150  
158  
166  
174  
182  
190  
198  
206  
214  
222  
230  
238  
246  
254  
262  
270  
278  
286  
294  
302  
310  
318  
326  
334  
342  
350  
358  
366  
374  
382  
390  
398  
406  
414  
422  
430  
438  
446  
454  
462  
470  
478  
486  
494  
502  
510  
518  
526  
534  
542  
550  
558  
566  
574  
582  
590  
598  
606  
614  
622  
630  
638  
646  
654  
662  
670  
678  
686  
694  
702  
710  
718  
726  
734  
742  
750  
758  
766  
774  
782  
790  
798  
806  
814  
822  
830  
838  
846  
854  
862  
870  
878  
886  
894  
902  
910  
918  
926  
934  
942  
950  
958  
966  
974  
982  
990  
998  
1006  
1014  
1022  
1030  
1038  
1046  
1054  
1062  
1070  
1078  
1086  
1094  
1102  
1110  
1118  
1126  
1134  
1142  
1150  
1158  
1166  
1174  
1182  
1190  
1198  
1206  
1214  
1222  
1230  
1238  
1246  
1254  
1262  
1270  
1278  
1286  
1294  
1302  
1310  
1318  
1326  
1334  
1342  
1350  
1358  
1366  
1374  
1382  
1390  
1398  
1406  
1414  
1422  
1430  
1438  
1446  
1454  
1462  
1470  
1478  
1486  
1494  
1502  
1510  
1518  
1526  
1534  
1542  
1550  
1558  
1566  
1574  
1582  
1590  
1598  
1606  
1614  
1622  
1630  
1638  
1646  
1654  
1662  
1670  
1678  
1686  
1694  
1702  
1710  
1718  
1726  
1734  
1742  
1750  
1758  
1766  
1774  
1782  
1790  
1798  
1806  
1814  
1822  
1830  
1838  
1846  
1854  
1862  
1870  
1878  
1886  
1894  
1902  
1910  
1918  
1926  
1934  
1942  
1950  
1958  
1966  
1974  
1982  
1990  
1998  
2006  
2014  
2022  
2030  
2038  
2046  
2054  
2062  
2070  
2078  
2086  
2094  
2102  
2110  
2118  
2126  
2134  
2142  
2150  
2158  
2166  
2174  
2182  
2190  
2198  
2206  
2214  
2222  
2230  
2238  
2246  
2254  
2262  
2270  
2278  
2286  
2294  
2302  
2310  
2318  
2326  
2334  
2342  
2350  
2358  
2366  
2374  
2382  
2390  
2398  
2406  
2414  
2422  
2430  
2438  
2446  
2454  
2462  
2470  
2478  
2486  
2494  
2502  
2510  
2518  
2526  
2534  
2542  
2550  
2558  
2566  
2574  
2582  
2590  
2598  
2606  
2614  
2622  
2630  
2638  
2646  
2654  
2662  
2670  
2678  
2686  
2694  
2702  
2710  
2718  
2726  
2734  
2742  
2750  
2758  
2766  
2774  
2782  
2790  
2798  
2806  
2814  
2822  
2830  
2838  
2846  
2854  
2862  
2870  
2878  
2886  
2894  
2902  
2910  
2918  
2926  
2934  
2942  
2950  
2958  
2966  
2974  
2982  
2990  
2998  
3006  
3014  
3022  
3030  
3038  
3046  
3054  
3062  
3070  
3078  
3086  
3094  
3102  
3110  
3118  
3126  
3134  
3142  
3150  
3158  
3166  
3174  
3182  
3190  
3198  
3206  
3214  
3222  
3230  
3238  
3246  
3254  
3262  
3270  
3278  
3286  
3294  
3302  
3310  
3318  
3326  
3334  
3342  
3350  
3358  
3366  
3374  
3382  
3390  
3398  
3406  
3414  
3422  
3430  
3438  
3446  
3454  
3462  
3470  
3478  
3486  
3494  
3502  
3510  
3518  
3526  
3534  
3542  
3550  
3558  
3566  
3574  
3582  
3590  
3598  
3606  
3614  
3622  
3630  
3638  
3646  
3654  
3662  
3670  
3678  
3686  
3694  
3702  
3710  
3718  
3726  
3734  
3742  
3750  
3758  
3766  
3774  
3782  
3790  
3798  
3806  
3814  
3822  
3830  
3838  
3846  
3854  
3862  
3870  
3878  
3886  
3894  
3902  
3910  
3918  
3926  
3934  
3942  
3950  
3958  
3966  
3974  
3982  
3990  
3998  
4006  
4014  
4022  
4030  
4038  
4046  
4054  
4062  
4070  
4078  
4086  
4094  
4102  
4110  
4118  
4126  
4134  
4142  
4150  
4158  
4166  
4174  
4182  
4190  
4198  
4206  
4214  
4222  
4230  
4238  
4246  
4254  
4262  
4270  
4278  
4286  
4294  
4302  
4310  
4318  
4326  
4334  
4342  
4350  
4358  
4366  
4374  
4382  
4390  
4398  
4406  
4414  
4422  
4430  
4438  
4446  
4454  
4462  
4470  
4478  
4486  
4494  
4502  
4510  
4518  
4526  
4534  
4542  
4550  
4558  
4566  
4574  
4582  
4590  
4598  
4606  
4614  
4622  
4630  
4638  
4646  
4654  
4662  
4670  
4678  
4686  
4694  
4702  
4710  
4718  
4726  
4734  
4742  
4750  
4758  
4766  
4774  
4782  
4790  
4798  
4806  
4814  
4822  
4830  
4838  
4846  
4854  
4862  
4870  
4878  
4886  
4894  
4902  
4910  
4918  
4926  
4934  
4942  
4950  
4958  
4966  
4974  
4982  
4990  
4998  
5006  
5014  
5022  
5030  
5038  
5046  
5054  
5062  
5070  
5078  
5086  
5094  
5102  
5110  
5118  
5126  
5134  
5142  
5150  
5158  
5166  
5174  
5182  
5190  
5198  
5206  
5214  
5222  
5230  
5238  
5246  
5254  
5262  
5270  
5278  
5286  
5294  
5302  
5310  
5318  
5326  
5334  
5342  
5350  
5358  
5366  
5374  
5382  
5390  
5398  
5406  
5414  
5422  
5430  
5438  
5446  
5454  
5462  
5470  
5478  
5486  
5494  
5502  
5510  
5518  
5526  
5534  
5542  
5550  
5558  
5566  
5574  
5582  
5590  
5598  
5606  
5614  
5622  
5630  
5638  
5646  
5654  
5662  
5670  
5678  
5686  
5694  
5702  
5710  
5718  
5726  
5734  
5742  
5750  
5758  
5766  
5774  
5782  
5790  
5798  
5806  
5814  
5822  
5830  
5838  
5846  
5854  
5862  
5870  
5878  
5886  
5894  
5902  
5910  
5918  
5926  
5934  
5942  
5950  
5958  
5966  
5974  
5982  
5990  
5998  
6006  
6014  
6022  
6030  
6038  
6046  
6054  
6062  
6070  
6078  
6086  
6094  
6102  
6110  
6118  
6126  
6134  
6142  
6150  
6158  
6166  
6174  
6182  
6190  
6198  
6206  
6214  
6222  
6230  
6238  
6246  
6254  
6262  
6270  
6278  
6286  
6294  
6302  
6310  
6318  
6326  
6334  
6342  
6350  
6358  
6366  
6374  
6382  
6390  
6398  
6406  
6414  
6422  
6430  
6438  
6446  
6454  
6462  
6470  
6478  
6486  
6494  
6502  
6510  
6518  
6526  
6534  
6542  
6550  
6558  
6566  
6574  
6582  
6590  
6598  
6606  
6614  
6622  
6630  
6638  
6646  
6654  
6662  
6670  
6678  
6686  
6694  
6702  
6710  
6718  
6726  
6734  
6742  
6750  
6758  
6766  
6774  
6782  
6790  
6798  
6806  
6814  
6822  
6830  
6838  
6846  
6854  
6862  
6870  
6878  
6886  
6894  
6902  
6910  
6918  
6926  
6934  
6942  
6950  
6958  
6966  
6974  
6982  
6990  
6998  
7006  
7014  
7022  
7030  
7038  
7046  
7054  
7062  
7070  
7078  
7086  
7094  
7102  
7110  
7118  
7126  
7134  
7142  
7150  
7158  
7166  
7174  
7182  
7190  
7198  
7206  
7214  
7222  
7230  
7238  
7246  
7254  
7262  
7270  
7278  
7286  
7294  
7302  
7310  
7318  
7326  
7334  
7342  
7350  
7358  
7366  
7374  
7382  
7390  
7398  
7406  
7414  
7422  
7430  
7438  
7446  
7454  
7462  
7470  
7478  
7486  
7494  
7502  
7510  
7518  
7526  
7534  
7542  
7550  
7558  
7566  
7574  
7582  
7590  
7598  
7606  
7614  
7622  
7630  
7638  
7646  
7654  
7662  
7670  
7678  
7686  
7694  
7702  
7710  
7718  
7726  
7734  
7742  
7750  
7758  
7766  
7774  
7782  
7790  
7798  
7806  
7814  
7822  
7830  
7838  
7846  
7854  
7862  
7870  
7878  
7886  
7894  
7902  
7910  
7918  
7926  
7934  
7942  
7950  
7958  
7966  
7974  
7982  
7990  
7998  
8006  
8014  
8022  
8030  
8038  
8046  
8054  
8062  
8070  
8078  
8086  
8094  
8102  
8110  
8118  
8126  
8134  
8142  
8150  
8158  
8166  
8174  
8182  
8190  
8198  
8206  
8214  
8222  
8230  
8238  
8246  
8254  
8262  
8270  
8278  
8286  
8294  
8302  
8310  
8318  
8326  
8334  
8342  
8350  
8358  
8366  
8374  
8382  
8390  
8398  
8406  
8414  
8422  
8430  
8438  
8446  
8454  
8462  
8470  
8478  
8486  
8494  
8502  
8510  
8518  
8526  
8534  
8542  
8550  
8558  
8566  
8574  
8582  
8590  
8598  
8606  
8614  
8622  
8630  
8638  
8646  
8654  
8662  
8670  
8678  
8686  
8694  
8702  
8710  
8718  
8726  
8734  
8742  
8750  
8758  
8766  
8774  
8782  
8790  
8798  
8806  
8814  
8822  
8830  
8838  
8846  
8854  
8862  
8870  
8878  
8886  
8894  
8902  
8910  
8918  
8926  
8934  
8942  
8950  
8958  
8966  
8974  
8982  
8990  
8998  
9006  
9014  
9022  
9030  
9038  
9046  
9054  
9062  
9070  
9078  
9086  
9094  
9102  
9110  
9118  
9126  
9134  
9142  
9150  
9158  
9166  
9174  
9182  
9190  
9198  
9206  
9214  
9222  
9230  
9238  
9246  
9254  
9262  
9270  
9278  
9286  
9294  
9302  
9310  
9318  
9326  
9334  
9342  
9350  
9358  
9366  
9374  
9382  
9390  
9398  
9406  
9414  
9422  
9430  
9438  
9446  
9454  
9462  
9470  
9478  
9486  
9494  
9502  
9510  
9518  
9526  
9534  
9542  
9550  
9558  
9566  
9574  
9582  
9590  
9598  
9606  
9614  
9622  
9630  
9638  
9646  
9654  
9662  
9670  
9678  
9686  
9694  
9702  
9710  
9718  
9726  
9734  
9742  
9750  
9758  
9766  
9774  
9782  
9790  
9798  
9806  
9814  
9822  
9830  
9838  
9846  
9854  
9862  
9870  
9878  
9886  
9894  
9902  
9910  
9918  
9926  
9934  
9942  
9950  
9958  
9966  
9974  
9982  
9990  
9998  
10006  
10014  
10022  
10030  
10038  
10046  
10054  
10062  
10070  
10078  
10086  
10094  
10102  
10110  
10118  
10126  
10134  
10142  
10150  
10158  
10166  
10174  
10182  
10190  
10198  
10206  
10214  
10222  
10230  
10238  
10246  
10254  
10262  
10270  
10278  
10286  
10294  
10302  
10310  
10318  
10326  
10334  
10342  
10350  
10358  
10366  
10374  
10382  
10390  
10398  
10406  
10414  
10422  
10430  
10438  
10446  
10454  
10462  
10470  
10478  
10486  
10494  
10502  
10510  
10518  
10526  
10534  
10542  
10550  
10558  
10566  
10574  
10582  
10590  
10598  
10606  
10614  
10622  
10630  
10638  
10646  
10654  
10662  
10670  
10678  
10686  
10694  
10702  
10710  
10718  
10726  
10734  
10742  
10750  
10758  
10766  
10774  
10782  
10790  
10798  
10806  
10814  
10822  
10830  
10838  
10846  
10854  
10862  
10870  
10878  
10886  
10894  
10902  
10910  
10918  
10926  
10934  
10942  
10950  
10958  
10966  
10974  
10982  
10990  
10998  
11006  
11014  
11022  
11030  
11038  
11046  
11054  
11062  
11070  
11078  
11086  
11094  
11102  
11110  
11118  
11126  
11134  
11142  
11150  
11158  
11166  
11174  
11182  
11190  
11198  
11206  
11214  
11222  
11230  
11238  
11246  
11254  
11262  
11270  
11278  
11286  
11294  
11302  
11310  
11318  
11326  
11334  
11342  
11350  
11358  
11366  
11374  
11382  
11390  
11398  
11406  
11414  
11422  
11430  
11438  
11446  
11454  
11462  
11470  
11478  
11486  
11494  
11502  
11510  
11518  
11526  
11534  
11542  
11550  
11558  
11566  
11574  
11582  
11590  
11598  
11606  
11614  
11622  
11630  
11638  
11646  
11654  
11662  
11670  
11678  
11686  
11694  
11702  
11710  
11718  
11726  
11734  
11742  
11750  
11758  
11766  
11774  
11782  
11790  
11798  
11806  
11814  
11822  
11830  
11838  
11846  
11854  
11862  
11870  
11878  
11886  
11894  
11902  
11910  
11918  
11926  
11934  
11942  
11950  
11958  
11966  
11974  
11982  
11990  
11998  
12006  
12014  
12022  
12030  
12038  
12046  
12054  
12062  
12070  
12078  
12086  
12094  
12102  
12110  
12118  
12126  
12134  
12142  
12150  
12158  
12166  
12174  
12182  
12190  
12198  
12206  
12214  
12222  
12230  
12238  
12246  
12254  
12262  
12270  
12278  
12286  
12294  
12302  
12310  
12318  
12326  
12334  
12342  
12350  
12358  
12366  
12374  
12382  
12390  
12398  
12406  
12414  
12422  
12430  
12438  
12446  
12454  
12462  
12470  
12478  
12486  
12494  
12502  
12510  
12518  
12526  
12534  
12542  
12550  
12558  
12566  
12574  
12582  
12590  
12598  
12606  
12614  
12622  
12630  
12638  
12646  
12654  
12662  
12670  
12678  
12686  
12694  
12702  
12710  
12718  
12726  
12734  
12742  
12750  
12758  
12766  
12774  
12782  
12790  
12798  
12806  
12814  
12822  
12830  
12838  
12846  
12854  
12862  
12870  
12878  
12886  
12894  
12902  
12910  
12918  
12926  
12934  
12942  
12950  
12958  
12966  
12974  
12982  
12990  
12998  
13006  
13014  
13022  
13030  
13038  
13046  
13054  
13062  
13070  
13078  
13086  
13094  
13102  
13110  
13118  
13126  
13134  
13142  
13150  
13158  
13166  
13174  
13182  
13190  
13198  
13206  
13214  
13222  
13230  
13238  
13246  
13254  
13262  
13270  
13278  
13286  
13294  
13302  
13310  
13318  
13326  
13334  
13342  
13350  
13358  
13366  
13374  
13382  
13390  
13398  
13406  
13414  
13422  
13430  
13438  
13446  
13454  
13462  
13470  
13478  
13486  
13494  
13502  
13510  
13518  
13526  
13534  
13542  
13550  
13558  
13566  
13574  
13582  
13590  
13598  
13606  
13614  
13622  
13630  
13638  
13646  
13654  
13662  
13670  
13678  
13686  
13694  
13702  
13710  
13718  
13726  
13734  
13742  
13750  
13758  
13766  
13774  
13782  
13790  
13798  
13806  
13814  
13822  
13830  
13838  
13846  
13854  
13862  
13870  
13878  
13886  
13894  
13902  
13910  
13918  
13926  
13934  
13942  
13950  
13958  
13966  
13974  
13982  
13990  
13998  
14006  
14014  
14022  
14030  
14038  
14046  
14054  
14062  
14070  
14078  
14086  
14094  
14102  
14110  
14118  
14126  
14134  
14142  
14150  
14158  
14166  
14174  
14182  
14190  
14198  
14206  
14214  
14222  
14230  
14238  
14246  
14254  
14262  
14270  
14278  
14286  
14294  
14302  
14310  
14318  
14326  
14334  
14342  
14350  
14358  
14366  
14374  
14382  
14390  
14398  
14406  
14414  
14422  
14430  
14438  
14446  
14454  
14462  
14470  
14478  
14486  
14494  
14502  
14510  
14518  
14526  
14534  
14542  
14550  
14558  
14566  
14574  
14582  
14590  
14598  
14606  
14614  
14622  
14630  
14638  
14646  
14654  
14662  
14670  
14678  
14686  
14694  
14702  
14710  
14718  
14726  
14734  
14742  
14750  
14758  
14766  
14774  
14782  
14790  
14798  
14806  
14814  
14822  
14830  
14838  
14846  
14854  
14862  
14870  
14878  
14886  
14894  
14902  
14910  
14918  
14926  
14934  
14942  
14950  
14958  
14966  
14974  
14982  
14990  
14998  
15006  
15014  
15022  
15030  
15038  
15046  
15054  
15062  
15070  
15078  
15086  
15094  
15102  
15110  
15118  
15126  
15134  
15142  
15150  
15158  
15166  
15174  
15182  
15190  
15198  
15206  
15214  
15222  
15230  
15238  
15246  
15254  
15262  
15270  
15278  
15286  
15294  
15302  
15310  
15318  
15326  
15334  
15342  
15350  
15358  
15366  
15374  
15382  
15390  
15398  
15406  
15414  
15422  
15430  
15438  
15446  
15454  
15462  
15470  
15478  
15486  
15494  
15502  
15510  
15518  
15526  
15534  
15542  
1555



FIGURE 2

1 ATGGAGGGCT GCATGGGGGA GGAGTCGTTT CAGATGTGGG AGCTCAATCG  
51 GCGCCTGGAG GCCTACCTGG GCCGGGTCAA GGCGCTGGAG GAGCAGAATG  
101 AGCTGCTCAG CGCCGGACTC GGGGGGCTCC GCGGACAATC CGCGGACAEC  
151 TCCTGGCGGG CGCATGCCGA CGACGAGCTG GCGGCCCTGC GTGCGCTCGT  
201 TGACCAACGC TGGCGGGAGA AGCACGCGGC CGAGGTGGCG CGCGACAACC  
251 TGGCTGAAGA GCTGGAGGGC GTGGCAGGCC GATGCGAGCA GCTGCGGCTG  
301 GCCCGGGAGC GGAOGACGGA GGAGGTAGCC CGCAACCGGC GCGCCGTCGA  
351 GGCAGAGAAA TGCGCCCGGG CCTGGCTGAG TAGCCAGGGG GCAGAGCTGG  
401 AGCGcGAGcT AGAGGCTcTA CGCGTgGCGC ACGAGGAGGA GCGCGTCGGT  
451 CTGAACGCGC AGgCtgcCTG TGCCCCCGCc cTGCCCGcGc cgcCCCGgGcc  
501 tccccgcgCCG GCCCCGGAGG tAGAGGAGcT GGcAAGGCGA CTGGGCGAGG  
551 CGTGGCGCGG GGCAGTGCGC GcCTACCAGG AGCGCGTGGC ACACATGGAG  
601 ACGTCGCTGG ACCAGAcCCG CGAGCGCCTG GCCCCGGCGG TGCAGGGTGC  
651 CCGCGAGGTC CGCCTGGAGC TACAGCAGCT CCAGGCTGAG CGCGGAGGCC  
701 TCCTGGAGCG CAGGGCAGCG TTGGAACAGA GGTGGAGGG CCGCTGgcag  
751 GaGcGGcTGC GGGcTACTGA AAAGTTCCAG CTGGCTGTGG AGGCCCTGGA  
801 GCAGGAGAAA CAGGGCCTAC AGAGCCAGAT CGCTCAGGTC CTGGAAGGTC  
851 GGCAGCAGCT GCGCACCTC AAGATGTCCC TCAGCCTGGA GGTGGCCACG  
901 TACAGGACCC TCCTGGAGGC TGAGAACTCC CGGCTGCAAA CACCTGGCGG  
951 TGGCTCCAAG ACTTCCCTCA GCTTTCAGGA CCCCAGCTG GAGCTGCAAT  
1001 TCCCTAGGAC CCCAGAGGGC CGGCGTCTTG GATCTTTGCT CCCAGTCCTG  
1051 AGCCCCAACTT CCCTCCCCTC ACCCTTGCCT GCTACCCTTG AGACACCTGT  
1101 GCCAGCCTTT CTTAAGAACC AAGAATTCTT CCAGGCCCGT ACCCCTACCT  
1151 TGGCCAGCAC CCCCATCCCC CCCACACCTC AGGCACCCTC TCCTGCTGTA  
1201 GATGEAGAGA TCAGAGCCCA GGATGCTCCT CTCTCTCTGC TCCAGACACA  
1251 GGGTGGGAGG AAACAGGCTC CAGAGCCCCT GCGGGCTGAA GCCAGGGTGG  
1301 CCATTCTGTC CAGCGTCCTG CCTGGACCAG AGGAGCCTGG GGGCCAGCGG  
1351 CAAGAGGCCA GTACAGGCCA GTCCCCAGAG GACCATGCCT CCTTGGCACC  
1401 ACCCCTCAGC CcTgACCACT CCAGTTTAGA GGCTAAGGAT GGAGAATCCG  
1451 GTGGGTCTAG AGTGTTCAGC ATATGCCGAG GGGAAAGGTGA AGGGCAAATC

FIGURE 2 (Continued)

1501 TGGGGGTTGG TAGAGAAAGA AACAGCCATA GAGGGCAAAG TGGTAAGCAG  
1551 CTTGCAGCAG GAAATATGGG AAGAAGAGGA TCTAAACAGG AAGGAAATCC  
1601 AGGACTCCCA GGTTCCTTTG GAAAAAGAAA CCCTGAAGTC TCTGGGAGAG  
1651 GAGATTCAAG AGTCACTGAA GACTCTGGAA AACCAGAGCC ATGAGACACT  
1701 AGAAAGGGAG AATCAAGAAT GTCCGAGGTC TTTAGAAGAA GACTTAGAAA  
1751 CACTAAAAAG TCTAGAAAAG GAAAATAAAA GAGCTATTAA AGGATGTGGA  
1801 GGTAGTGAGA CCTCTAGAAA AAGAGGCTGT AGGCAACTTA AGCCTACAGG  
1851 AAAAGAGGAC ACACAGACAT TGCAATCCCT GCAAAGGAG AATCAAGAAC  
1901 TAATGAAATC TCTTGAAGGT AATCTAGAGA CATTTTATT TCCAGGAACG  
1951 GAAATCAAG AATTAGTAAG TTCTCTGCAA GAGAACTTAG AGTCATTGAC  
2001 AGCTCTGGAA AAGGAGAATC AAGAGCCACT GAGATCTCCA GAAGTAGGGG  
2051 ATGAGGAGGC ACTGAGACCT CTGACAAAGG AGAATCAGGA ACCCCTGAGG  
2101 TCTCTTGAAG ATGAGAACAA AGAGGCCTTT AGATCTCTAG AAAAAGAGAA  
2151 CCAGGAGCCA CTGAAGACTC TAGAAGAAGA GGACCAGAGT ATTGTGAGAC  
2201 CTCTAGAAAC AGAGAATCAC AAATCACTGA GGTCTTTAGA AGAACAGGAC  
2251 CAAGAGACAT TGAGAACTCT TGAAAAGAG ACTCAACAGC GACGGAGGTC  
2301 TCTAGGGGAA CAGGATCAGA TGACATTAAG ACCCCCAGAA AAAGTGGATC  
2351 TAGAACCACT GAAGTCTCTT GACCAGGAGA TAGCTAGACC TCTTGAAAAT  
2401 GAGAATCAAG AGTTCTTAAA GTCACTCAA GAAGAGAGCG TAGAGGCAGT  
2451 AAAATCTTTA GAAACAGAGA TCCTAGAATC ACTGAAGTCT GCGGGACAAG  
2501 AGAACCTGGA AACACTGAAA TCTCCAGAAA CTCAAGCACC ACTGTGGACT  
2551 CCAGAAGAAA TAAATAAATC AGGGGGCAAT GAATCCTCTA GAAAAGGAAA  
2601 TTCAAGAACC ACTGGAGTCT GTGGAAGTGA ACCAAGAGAC ATTCAGACTC  
2651 CTGGAAGAGG AGAATCAGGA ATCATTGAGA TCTCTGGGAG CATGGAACCT  
2701 GGAGAATTTG AGATCTCCAG AGGAGTAGAC AAGGAAAGTC AAAGGAATCT  
2751 GGAAGAGGAA GAGAACCTGG GAAAGGGAGA GTACCAAGAG TCACTGAGGT  
2801 CTCTGGAGGA GGAGGGACAG GAGCTGCCGC AGTCTGCAGA TGTGCAGAGG  
2851 TGGAAGATA CGGTGGAGAA GGACCAAGAA CTGGCTCAGG AAAGCCCTCC

FIGURE 2 (Continued)

2901 TGGGATGGCT GGAGTGGAAA ATAAgGatga Ggcagagctg AatctAaGGg  
2951 agcaGgatGG CTTCACTGGG AAGGAGGAGG TGGTAGAGCA GGGAGAGCTG  
3001 AATGCCACAG AGGAGGTCTG GTTCCCAGGC GAGGGGCACC CAGAGAACCC  
3051 TGAGCCCCAA GAGCAGAGAG GCCTGGTTGA GGGAGCCAGT gTGAAGGGAg  
3101 GGgctgagGg cctCcaggac CctgaAgGgC AATCACAACA GGTGGGGACc  
3151 CCAGGcCTCC AGGCTCCCCA GGGgctgCca gaGgagatag agCCcctGgt  
3201 GgAagatgaT GTGGCCcCAG GGGGTGACca AGCCTCCCCA GAGGTCATGT  
3251 TGGGGTCAGA GCCTGCCATG GGTGAGTCTG CTGCGGGagC TGAGCCAGgc  
3301 ctGGGgcaGG GGgtGGgaGG gCtGGGGgaC CcaggCcATC TgACCAGGGA  
3351 AGAGGTGATG GAACCACCCC TGGAAGAGGA GAGTTTGGAG GCAAAGAGGG  
3401 TTCagGGCTT GGAAGGGCCT AGAAAGGACC TAGAGGAGGC AGGTGGTCTG  
3451 GGGACAGAGT TCTCCGAGCT GCCTGGGAAG AGCAGAGACC CTTGGgAGCC  
3501 TCCCAGGGAG GGTAGGGAGG AGTCagAGGC TGAGGCCCCC AGGGGAGCAG  
3551 AGGAGGCGTT CCCTGCTGAG ACCCTGGGCC AACTTGGAAG TGATGCCCCCT  
3601 TCACCTTGGC CTCTGGGGTC AGAGGAAGCT GAGGAGGATG TACCACCAGT  
3651 GCTGGTCTCC CCCAGCCCAA CGTACACCCC GATCCTGGAA GATGCCCCTG  
3701 GGCTTcAGCC TCAGGCTGAA GGGAGTCAGG AGGCTAGCTG GGGGGTGCAG  
3751 GGGAGGGCTG AAGCTGGGAA AGTAGAGAGC GAGCAGGAGG AGTTGGGTTC  
3801 TGGGGAGATC CCGAGGGGCC TCCAGGAGGA AGGGGAGGAG AGCAGAGAAG  
3851 AGAGCGAGGA GGATGAGCTC GGGGAGACCC TTCCAGACTC CACTCCCCTG  
3901 GGCTTCTACC TCAGGTCCCC CACCTCCCCC AGGTGGACCC CACTGGAGAG  
3951 CAGAGGCCAC CCCCTCAAGG AGACTGGAAA GGAGGGCTGG GATCCTGCTG  
4001 TCCTGGCTTC CGAGGGCCTT GAGGAiCCCT CAGAAAAGGA GGAGGGGGAG  
4051 GAGGGAGAAG AGGAGTGTGG CCGTGA CTCT GACCTGTCAG AAGAATTGA  
4101 GGACCTGGGG ACTGAGGCAC CTTTTCTTCC TGGGGTCCCT GGGGAGGTGG

FIGURE 2 (Continued)

4151 CAGAACCTCT GGGCCAGgtg cCcCaGCTGC TACTGGATCC TGCAGCCTGG  
4201 GATCGAGATG GGGAGTCTGA TGGGTTTGCA GATGAGGAag AAAGTGGGGA  
4251 GGAGGGAGAG GAGGATCAGG AGGAGGGGAG GGAGCCAGGG GCTGGGCGGT  
4301 GGGGGCCAGG GTCTTCTGTT GGCAGCCTCC AGGCCCTGAG TAGCTCCAG  
4351 AGAGGGGAAT TCCTGGAGTC TGATTCTGTA AGTGTGACGG TCCCCTGGGA  
4401 TGACAGCTTG AGGGGTGCAG TGGCTGGTGC CCCCAAGACT GCCCTGGAAA  
4451 CGGAGTCCCA GGACAGTGCT GAGCCTTCTG GCTCAGAGGA AGAGTCTGAC  
4501 CCTGTTTCCT TGGAGAGGGA GGACAAAGTC CCTGGCCCTC TAGAGATCCC  
4551 CAGTGGGATG GAGGATGCAG GCCCAGGGGC AGACATCATT GGTGTTAATG  
4601 GCCAGGGTCC CAACTTGGAG GGGAAAGTCAC AGCATGTAAA TGGGGGAGTA  
4651 ATGAACGGGC TGGAGCAGTC TGAGGAAAGT GGGGCAAGGA ATGCGCTAGT  
4701 CTCTGAGGGA GACCGAGGGA GCCCCTTTCA GGAGGAGGAG GGGAGTGCTC  
4751 TGAAGAGGTC TTCGGCAGGG GCTCCTGTTC ACCTGGGCCA GGGTCAGTTC  
4801 CTGAAGTTCA CTCAGAGGGA AGGAGATAGA GAGTCCTGGT CCTCAGGGGA  
4851 GGAC

FIGURE 3

1 MEGCMGEESF QMWELNRRLE AYLGRVKALE EQNELLSAGL GGLRRQSADT  
51 SWRAHADDEL AALRALVDQR WREKHAAEVA RDNLAEELEG VAGRCEQLRL  
101 ARERTTEEVA RNRRAVEAEK CARAWLSSQG AELEEREAL RVAHEEERVG  
151 LNAQAACAPR LPAPPRPPAP APEVEELARR LGEAWRGAVR GYQERVAHME  
201 TSLDOTRERL ARAVOGAREV RLELQQLQAE RGGLLERRAA LEQRLGRWQ  
251 ERLRATEKFO LAVEALEQEK QGLQSQIAQV LEGROQLAHL KMSLSLEVAT  
301 YRTLLEAENS RLOTGGGSK TSLSFQDPKL ELQFPRTPEG RRLGSLLPVL  
351 SPTSLSPLP ATLETPVPAF LKNQEFLOAR TPTLASTPIP PTPQAPSPAV  
401 DAEIRAQDAP LSLLOTGGR KOAPEPLRAE ARVAIPASVL PGPEEPGGQR  
451 QEASTGOSPE DHASLAPPLS PDHSSLEAKD GESGGSRVFS ICRGEGEGQI  
501 WGLVEKETAI EGKVSSLOQ EIWEEEDLNR KEIQDSQVPL EKETLKSIGE  
551 EIQESLKTLE NQSHETLERE NOECPRSLEE DLETLSLEK ENKRAIKCGG  
601 GSETSRKRC RQKPTGKED TOTLQSLQKE NQELMKSLEG NLETFLFPGT  
651 ENQELVSSLO ENLESITALE KENQEPLRSP EVGDEEALRP LTKENQEPLR  
701 SLEDENKEAF RSLEKENQEP LKTLLEEDQS IVRPLETENH KSLRSLEEQD  
751 QETLRITLKE TOORRRSLGE QDOMTLRPPE KVDLEPLKSL DQEIARPLEN  
801 ENQEFKSLK EESVEAVKSL ETEILESLSKS AGQENLETLK SPETQAPLWT  
851 PEEINKSGGN ESSRKGNRT TGVCGSEPRD IQTPGRGESG IIEISGSMEP  
901 GEFEISRGVD KESQNLLEE ENLGKGEYQE SLRSLEEEO ELPOSADVQR  
951 WEDTVEKDQE LAQESPPGMA GVENKDEAEL NLREQDGFTG KEEVVEQGE  
1001 NATEEVWFFG BHPENPEPK EQRGLVEGAS VKGGAEGLOD PEGOSQOVGT  
1051 PGLQAPQGLP EATEPLVEDD VAPGGDOASP EVMLGSEPAM GESAAGAEPG

FIGURE 3 (Continued)

1101 LGQGVGGLGD PGHLTREEVM EPPLEESLE AKRVQGLEGP RKDLEEAGGL  
1151 GTEFSELP GK SRDPWEPPRE GRESEAEAP RGAEEAFPAE TLGHTGSDAP  
1201 SPWPLGSEEA EEDVPPVLVS PSPTYTPILE DAPGLQPQAE GSQEASWGVQ  
1251 GRAEAGKVES EQEELGSGEI PEGLOEEGEE SREESEEDL GETLPDSTPL  
1301 GFYLRSP TSP RWTPLSRGH PLKETGKEGW DPAVLASEGL EEPSEKEEGE  
1351 EGEEECGRDS DLSEEFEDLG TEAPFLPGVP GEVAEPLGQV PQLLLDPAAW  
1401 DRDGESDGFA DEESGEEGE EDQEEGREPG AGRWGPSSV GSLOALSSSQ  
1451 RGEFLES DSV SVSVPWDDSL RGAVAGAPKT ALETESQDSA EPSGSEESD  
1501 FVSLEREDKV PGPLEIPSGM EDAGPGADII GVNGQGNLE GKSQHVNGGV  
1551 MNGLEQSEES GARNALVSEG DRGSPFOEEE GSALKRSSAG APVHLGGQGF -  
1601 LKFTQREGDR ESWSSGED